

Cryptic species of *Didymobothrium rudolphii* (Cestoda: Spathebothriidea) from the sand sole, *Solea lascaris*, off the Portuguese coast, with an analysis of their molecules, morphology, ultrastructure and phylogeny

J. F. MARQUES^{1*}, M. J. SANTOS², D. I. GIBSON³, H. N. CABRAL¹ and P. D. OLSON³

¹ Universidade de Lisboa, Faculdade de Ciências, Instituto de Oceanografia, Campo Grande, 1749-016 Lisboa, Portugal

² Universidade do Porto, Faculdade de Ciências, Departamento de Zoologia e Antropologia, Praça Gomes Teixeira, 4099-002 Porto and CIMAR/CIIMAR – Centro Interdisciplinar de Investigação Marinha e Ambiental, Rua dos Bragas, 177, 4050-123 Porto, Portugal

³ Department of Zoology, Natural History Museum, Cromwell Road, London SW7 5BD, UK

(Received 3 November 2006; revised 9 January 2007; accepted 10 January 2007; first published online 28 February 2007)

SUMMARY

Didymobothrium rudolphii (Cestoda: Spathebothriidea) was collected seasonally from the sand sole, *Solea lascaris*, off the northern, central and southern areas of the Portuguese coast. Morphological and molecular analyses were conducted in order to examine the possible existence of cryptic species and to facilitate the circumscription of their morphological boundaries. Data were compared between *D. rudolphii* specimens from each of the 3 geographical areas and 4 seasons, and principal components analysis of 18 morphological characters was used to detect differences. Two distinct genotypes were present with sequence divergences of 1.9% and 2.1% in the large subunit (lsrDNA) and second internal transcribed spacer (ITS-2) of ribosomal DNA (rDNA), respectively. The less common ‘central’ genotype was present only off the central area from summer to winter, whereas the ‘common’ genotype was present throughout the year off the northern and southern areas, but only during spring in the central area. No sequence variation was found within each genotype. The presence of 2 distinct genetic entities was supported by morphological analyses, which showed the ‘central’ genotype specimens to be more slender and elongate, although morphometric ranges overlapped considerably for most characters. Molecular phylogenetic analysis of 4 of the 5 known genera of the Spathebothriidea showed *Spathebothrium* to be the earliest branching lineage and the 2 genotypes of *Didymobothrium* formed a sister group to *Cyathocephalus*. The concordance of genetic differences with variation in host diet according to season and locality could account for sympatric speciation occurring in the central region of the Portuguese coast.

Key words: *Didymobothrium*, Spathebothriidea, large-subunit rDNA, ITS-2, cryptic speciation, systematics, *Solea lascaris*, *Bothrimonus*, *Cyathocephalus*, *Diplocotyle*, *Spathebothrium*, Portuguese coast.

INTRODUCTION

Tapeworms of the order Spathebothriidea are unique in exhibiting serially repeated reproductive organs without the accompanying segmentation that is the hallmark of the Cestoda. They comprise only 5 genera, are widely but disjunctively spread across the northern hemisphere, and have a similarly disparate pattern of host associations, occurring in a small number of marine, euryhaline and freshwater teleosts and chondrosteans (Gibson, 1994). In addition, ovigerous adults of both freshwater and marine forms are known to develop in Amphipoda, circumventing the necessity of a vertebrate final

host in the life-cycle (Gibson and Valtonen, 1983; Protasova and Roytman, 1995; Davydov *et al.* 1997; Okaka, 2000; Mackiewicz, 2003; Poddubnaya *et al.* 2003, 2005). Taken together, such characteristics suggest that the Spathebothriidea is a relict group of tapeworms whose extant forms represent refugial lineages of a clade that was once more diverse. Moreover, molecular phylogenetic studies indicate that the Spathebothriidea does indeed form a basal lineage of the Eucestoda (possibly the sister group to the other Eucestoda; for a review see Olson and Tkach, 2005), from which it is inferred that they exhibit the ancestral condition of the unsegmented Cestodaria (i.e. Amphilinidea and Gyrocotylidea) in combination with the derived condition of having proglottides (i.e. exhibiting serially-repeated reproductive organs). They may therefore represent the key step in the evolution of the reproductive strategy that now characterizes the vast majority of

* Corresponding author: Universidade de Lisboa, Faculdade de Ciências, Instituto de Oceanografia, Campo Grande, 1749-016 Lisboa, Portugal. Tel: +351 217 500 826. Fax: +351 217 500 207. E-mail: jimarques@fc.ul.pt

cestodes. Nevertheless, they are rarely reported and remain largely unknown in most aspects of their biology.

The systematics and affinities of the Spathebothriidea with other cestode orders have been controversial (e.g. Burt and Sandeman, 1969; Gibson, 1994; Brunanska *et al.* 2005; Poddubnaya *et al.* 2005), as has the validity of 3 of the 5 genera: *Bothrimonus* Duvernoy, 1842, *Diplocotyle* Krabbe, 1874 and *Didymobothrium* Nybelin, 1922. Burt and Sandeman (1969) presented a detailed review of the systematics and morphology of these genera, concluding that the differences used to separate them were not valid and that *Diplocotyle* and *Didymobothrium* should therefore be regarded as junior synonyms of *Bothrimonus*. Gibson (1994), following Nybelin (1922), differentiated *Bothrimonus* from the other two genera by the lack of a septum dividing the lumina of the bothridia, and *Diplocotyle* and *Didymobothrium* from one another by the arrangement of the genital pores: ventral in the former and irregularly alternating in the latter. Nevertheless, at about the same time, Protasova and Roytman (1995) considered *Didymobothrium* a *nomen dubium*, suggesting that *D. rudolphii* (Monticelli, 1890) Nybelin, 1922 might prove to be a synonym of *Diplocotyle olrikii* Krabbe, 1874. Detailed morphological studies have been conducted on *Diplocotyle nylandica* (Schneider, 1902) (see MacKinnon and Burt, 1984) and *D. olrikii* (see Brunanska *et al.* 2005; Poddubnaya *et al.* 2005), but *Bothrimonus* spp. (*sensu stricto*) and *Didymobothrium rudolphii* remain poorly known, only the ultrastructure of the vitelline system of the latter having been studied in detail by Poddubnaya *et al.* (2006).

In a parasitological survey of the soles *Solea lascaris* (Risso, 1810) and *S. impar* Bennett, 1831 (Pisces, Pleuronectiformes) conducted along the northeast Atlantic coast and in the Mediterranean (including a sampling site near Lisbon, on the Portuguese coast), Renaud and Gabrion (1988) found these species to be infected with what they considered *Bothrimonus nylandicus*, following the work of Burt and Sandeman (1969). Using allozyme electrophoresis, they demonstrated 2 cryptic species occurring in both the Atlantic and the Mediterranean. Whereas one of the cryptic species was found throughout the year, the other first appeared in spring and was absent by the end of the summer. To account for this, the authors' formulated a number of hypotheses regarding the longevity of the different 'species' and the potential for different intermediate host specificity driving speciation.

Off the coast of Portugal, *D. rudolphii* is commonly found infecting *S. lascaris* throughout the year, enabling continual sampling of the parasite populations from this commercially important host species. Although Renaud and Gabrion (1988) reported




Fig. 1. Sampling areas off the Portuguese coast.

D. rudolphii (as *B. nylandicus*) to infect *S. impar* off the Portuguese coast, the taxonomic validity of this host is currently in question (Infante *et al.* 2004), and no specimen sampled in the present study corresponded to the description of *S. impar*. In order to investigate further the presence of cryptic species and to attempt to circumscribe the morphometric and molecular variation exhibited by these entities, parasites were collected seasonally from the northern, central and southern areas off the Portuguese coast (Fig. 1). This material also provided an opportunity to study their morphology in detail for the first time, including ultrastructural and histological features, thus significantly augmenting previous accounts published more than 80 years ago (i.e. Monticelli, 1890, 1892; Nybelin, 1922).

MATERIALS AND METHODS

Specimen collection and vouchers

During 2003, 480 specimens of the sand sole, *Solea lascaris*, were obtained from commercial fishing vessels operating off the northern ($n=158$), central ($n=169$) and southern ($n=153$) areas of the Portuguese coast (Fig. 1) and examined for the

presence of *D. rudolphii*. Cestode specimens were fixed in 10% neutral-buffered formalin, 2.5% phosphate-buffered glutaraldehyde or 100% analytical grade ethanol, depending on the technique to be used subsequently. The stomach contents of the fish were also examined, in order to gain insights into host feeding ecology and its relationship to infection patterns. Voucher specimens, including whole-mounted and sectioned material used for morphological study, were deposited in the Museu Nacional de História Natural (Museu Bocage), Lisbon (MNHN MB8-19), and whole-mounted, partial specimens of 12 of the 24 samples used for molecular analysis were deposited in the Natural History Museum, London (BMNH 2006.10.4.9-20).

Molecular analyses

Eight individuals from each of the 3 sampling areas (2 per season; 24 in total) were fixed in pure ethanol for molecular analysis. Ethanol was replaced with Tris-ethylenediamine tetracetic acid (EDTA) buffer (pH 8.0) by soaking overnight, and genomic DNA was extracted using a Qiagen DNeasyTM tissue kit, with modifications as described by Agustí *et al.* (2005). Two nuclear ribosomal regions were characterized: the large-subunit (lsrDNA) which has proven to be informative for both phylogenetic and diagnostic studies of the Cestoda (e.g. Brickle *et al.* 2001; Olson *et al.* 2001; Reyda and Olson, 2003; Agustí *et al.* 2005; Aznar *et al.* 2007) and the second internal transcribed spacer (ITS-2) which has been commonly used for investigating inter- and intraspecific variation in flatworms (e.g. Verneau *et al.* 1997; Zehnder and de Chambrier, 2000; Olson *et al.* 2002; see Olson and Tkach, 2005 and Nolan and Cribb, 2005). Partial lsrDNA (D1-D3; ~1400 bp) and complete ITS-2 (~600 bp) genes were amplified by polymerase chain reaction (PCR) using oligonucleotide primers LSU5 (5' TAG GTC GAC CCG CTG AAY TTA AGC 3') + 1200R (5' GCA TAG TTC ACC ATC TTT CGG 3'), and ITS2-3S (5' GGT ACC GGT GGA TCA CGT GGC TAG TG 3') + ITS2-2 (5' CCT GGT TAG TTT CTT TTC CTC CGC 3'), respectively. Cycle sequencing was performed bidirectionally using the PCR primers and internal primers 300F (5' CAA GTA CCG TGA GGG AAA GTT 3') and ECD2 (5' CTT GGT CCG TGT TTC AAG ACG GG 3') in the case of the lsrDNA. Contiguous sequences were assembled and edited using SequencherTM (GeneCodes Corp.), screened for contamination via BLASTn (McGinnis and Madden, 2004) and deposited in GenBank under Accession numbers EF042920- EF042965.

To facilitate comparative sequence and phylogenetic analyses, collection and/or sequencing of additional spathebothriidean taxa was carried out

as follows: a gravid (progenetic) specimen of *Diplocotyle olrikii* was collected from *Gammarus* sp. (Amphipoda) at St Andrews, Scotland and both the lsrDNA and ITS-2 characterized; the ITS-2 was also characterized for *Spathebothrium simplex* Linton, 1922 and *Cyathocephalus truncatus* (Pallas, 1781) in order to complement lsrDNA sequences characterized previously for these taxa (see Olson *et al.* 2001).

In order to maximize the number of alignable characters, and thus provide the best estimates of relative divergences from the available sequences, data sets were constructed for the lsrDNA and ITS-2 representing only the spathebothriidean taxa and were thus analysed phylogenetically as unrooted networks. In addition, a third data set based on lsrDNA was constructed that included sequences representing 2 gyrocotylidean outgroups in order to root the spathebothriidean clade; of the ~1400 bp of lsrDNA characterized, only 881 (~63%) were alignable together with the gyrocotylidean sequences, whereas 1275 (~91%) were alignable when considering only the spathebothriidean taxa. One representative sequence of each of the 2 distinct *Didymobothrium* genotypes found were used in all analyses. Alignments were made by eye using the program MacClade 4.08 (Maddison and Maddison, 2005). Comparative and bootstrap analyses were performed using PAUP* 4.0b10 (Swofford, 2001), and phylogenetic affinities were estimated by Bayesian inference using MrBayes 3.1.2 (Ronquist and Huelsenbeck, 2003). Nucleotide substitution models were estimated for each data set individually using MrModeltest 1.1b (Nylander, 2004), a simplified version of ModelTest (Posada and Crandall, 1998) and a general time-reversible model of nucleotide substitution incorporating gamma-distributed among site rate variation (GTR + G for the ITS-2 data) or invariant sites (GTR + I for the lsrDNA data), or both (GTR + I + G for the lsrDNA data including the gyrocotylidean outgroup taxa), was specified and the analyses run over 0.5 million generations, sampling topologies every 100th generation. Other program parameters were as specified by Olson *et al.* (2003). Consensus trees were constructed using the 'sumt' command with 'burnin' = 20 and 'contype' = allcompat.

Morphometric analyses

For comparative morphometric analysis, 60 complete individuals (5 per season per locality) presenting different degrees of maturation were fixed in 70% ethanol, stained with iron acetocarmine (following Georgiev *et al.* 1986), dehydrated in an ethanol series, cleared with clove oil, mounted in Canada balsam and observed using light microscopy. Specimens were classified as 'immature' (without fully-formed reproductive organs), 'mature' (with

fully-formed reproductive organs, but without eggs) or 'gravid' (with fully-formed reproductive organs and with eggs) following Poddubnaya *et al.* (2003). Sixteen continuous and 2 meristic characters (Table 2) were measured using a compound microscope linked to image analysis software (Axiovision 3.1). Characters were chosen based on their diagnostic value following previous studies (Burt and Sandeman, 1969; Gibson, 1994; Pertierra, 2002; Hanzelová *et al.* 2005) and the original description (Monticelli, 1890). Since measurements of the reproductive organs vary according to the width of the individual, the width of the uterus, ovary and vitellarium were divided by the total width of the body in order to establish a ratio, expressed as a percentage of the body width, which could be compared meaningfully among the specimens. The number of testes was determined in mature proglottides only, since they degenerate as the worms become gravid. Unless otherwise indicated, measurements are presented in micrometres (μm) and given in the text as the mean or as a range. Only complete and either mature or gravid specimens were considered in the statistical analyses, and an average of the reproductive characters from 3 proglottides per individual was made. Following the results of genetic analyses, the morphometric data were pooled separately between the specimens originating in the north and south during all seasons, and for the central area from winter to summer. Metric data from the 2 genotypes (see molecular results) were compared with each other as well as with those published for *Diplocotyle* and '*Bothriomonus*' spp. *sensu lato* (Table 2). Differences between measurements of the 2 genotypes were tested at the significance level of 0.05 through an independent samples *t*-test (continuous variables) or Mann-Whitney (meristic variables) using the software package SPSS 13.0 (SPSS Inc., 2004).

Principal component analysis (PCA) was conducted in CANOCO 4.5 (ter Braak and Smilauer, 2002) to determine the multivariate relationship among the 16 continuous morphometric variables and to identify the most important of these for distinguishing potentially cryptic species. In order to have a better understanding of the variables influencing morphological variation, specimens from the central area collected during spring (which, according to the molecular results, exhibited the 'common' genotype) were excluded from this analysis. Variables showing the highest influence on the pattern found were tested for significant differences, at a level of 0.05, according to locality or season using the Kruskal-Wallis test carried out in SPSS 13.0. This non-parametric test was chosen as it allows the detection of differences in the distribution of values, determining if the tested samples come from the same population.

Scanning electron microscopy

Samples collected in the northern area during the summer of 2005 were processed for scanning electron microscopy (SEM). Specimens ($n=12$) were fixed in 2.5% phosphate-buffered glutaraldehyde, dehydrated in an ethanol series, critical-point dried in CO_2 , mounted on specimen stubs using a fine coating of Araldite (i.e. epoxy) glue, sputter coated with gold-palladium to a thickness of 20–40 nm and examined at 5 kV. To examine their internal anatomy by SEM, specimens in 100% ethanol were cut transversally or held on their side between glass slides and cut longitudinally using a razor blade. Other specimens were freeze-fractured in liquid nitrogen after first being infiltrated with epon and placed in a gelatine capsule to help stabilize the specimens during the freeze-fracture process. Cut or fractured specimens were then prepared for SEM.

Histology

Three individuals exhibiting varying degrees of maturation, also collected in the northern region during the summer of 2005, were used for histological sectioning in order to elucidate the arrangement and development of the reproductive organs and eggs. Individuals were fixed in Bouin's fixative, embedded in liquid paraffin, sectioned at 5 μm with a rotary microtome and stained with either haematoxylin and eosin or Periodic Acid Schiff (PAS) reagent. Specimens were then mounted in EntellanTM (Merck) and observed using light microscopy.

RESULTS

Comparative sequence analysis

Totals of 21 lsrDNA and 22 ITS-2 sequences were obtained from the 24 individuals analysed; specimens collected in the south during summer (i.e. during the period with the highest temperatures) usually proved the least amenable to genetic analysis. Comparative sequence analysis showed 2 distinct genotypes for both the lsrDNA and ITS-2. In both cases, the more common genotype (hereafter designated 'common') was found throughout the year in the north and south, and only during spring in the central region, whereas the less common genotype (hereafter designated 'central') was found only in the central region from summer to winter. Within each genotype, no sequence variation was observed for either rDNA region.

Comparison of the lsrDNA sequences (1301 nucleotides) showed 18 transitional and 7 transversional substitutions ($\text{ts}/\text{tv}=2.57$), or a raw genetic distance of 1.9%. Comparison of the ITS-2 sequences (612 nucleotides) showed 10 transitional and 3 transversional changes ($\text{ts}/\text{tv}=3.33$), or a raw genetic distance of 2.1%. Comparisons of sequence

Table 1. Comparison of genetic distances (% difference) between spathebothriidean taxa using large subunit rDNA (lsrDNA) and internal transcribed spacer 2 (ITS-2)

(Dip, *Diplocotyle orikii*; Cyt, *Cyathocephalus truncatus*; Dib:Com, *Didymobothrium rudolphii* 'common' genotype; Dib:Cen, *Didymobothrium rudolphii* 'central' genotype; Sps, *Spathebothrium simplex*.)

Taxon§	lsrDNA (N = 1275 bps)				ITS-2 (N = 588 bps)			
	Sps	Dip	Cyt	Dib:Com	Sps	Dip	Cyt	Dib:Com
Dip	5.6				6.1			
Cyt	6.1	4.2			7.7	4.1		
Dib:Com	7.5	7.1	6.0		8.7	5.4	6.9	
Dib:Cen	6.4	6.3	5.3	2.0	9.5	5.9	7.4	2.2




Fig. 2. Phylogenetic analyses of lsrDNA and ITS-2. (A) Rooted phylogram with relative branch lengths based on Bayesian inference (solid lines) and parsimony-based bootstrap analysis (dashed lines) of 888 bps of lsrDNA. (B) Unrooted phylograms based on 1275 bps of lsrDNA, and (C) on 588 bps ITS-2. Nodal support shown as posterior probabilities/bootstrap consensus (bold). GenBank sequence Accession numbers are shown parenthetically; those characterized in the present study are marked with an asterisk.

divergence between the 2 genotypes and other spathebothriidean genera showed *Cyathocephalus* to be most similar, with a raw distance of 5.3% from the central genotype and 6% from the common genotype. Other distances estimated are given in Table 1.

Phylogenetic analyses

Phylogenetic analysis of the lsrDNA data, including those of gyrocotylidean outgroup taxa (Fig. 2A), showed *Spathebothrium* to be the sister of the other spathebothriidean taxa and that the common and central genotypes of *Didymobothrium* formed a clade with *Cyathocephalus* and represent its sister lineage. Although both Bayesian and parsimony-based bootstrap analyses supported *Spathebothrium* as the earliest diverging taxon, slight differences were seen in the interrelationships of the other

spathebothriidean taxa, with nodal support being weak in both analyses. In the unrooted, individual analyses of lsrDNA and ITS-2 data sets (Fig. 2B and C), the 2 genotypes of *Didymobothrium* formed sister lineages with a high degree of divergence between them and, similarly, terminal branch lengths were long relative to the internal branch separating *Spathebothrium* + *Diplocotyle* from *Cyathocephalus* + the 2 *Didymobothrium* genotypes. Comparison of relative branch lengths between lsrDNA (Fig. 2B) and ITS-2 (Fig. 2C) showed lsrDNA data estimating slightly more equitable divergences between the lineages.

Morphological analyses

Morphometric comparison of the 'common' and 'central' genotypes of *D. rudolphii* as defined by the

Table 2. Morphometric comparison of the ‘common’ and ‘central’ genotypes of *Didymobothrium rudolphii* and other species of the Acrobothriidae as available in the literature

(Values correspond to the mean ± standard deviation and range (in parentheses) in µm (except where noted). * indicates significant differences between *D. rudolphii* genotypes ($P < 0.05$). §TL, total length; MW, maximum width; TLsclx, total length of the scolex; MWsclx, maximum width of the scolex; MTsclx, maximum thickness of the scolex; DistPore, distance between two consecutive female genital pores; DistCirr, distance between two consecutive cirri; MDCirr, maximum diameter of the cirrus; Nprogs, number of proglottides; MDtestes, maximum diameter of testes; Ntestes, number of testes; MLegg, maximum length of eggs; MWegg, maximum width of eggs; MLut, maximum length of the uterus; MWut, maximum width of the uterus; MLov, maximum length of the ovary; MWov, maximum width of the ovary; MWvt, maximum width of the vitellarium; Prout, percentage of strobila width occupied by the uterus; Proov, percentage of strobila width occupied by the ovary; Provt, percentage of strobila width occupied by vitellaria.

(1) as ‘*Bothrimonus nylandicus*’ from Schneider, 1902

(2) Adapted from Burt and Sandeman, 1969

(3) Nybelin, 1922

(4) Renaud and Gabrion, 1988.)

Variable§	<i>Didymobothrium rudolphii</i> ‘common genotype’ N=39	<i>Didymobothrium rudolphii</i> ‘central genotype’ N=14	<i>Diplocotyle nylandica</i> (1)	<i>Diplocotyle olrikii</i> (2)	<i>Bothrimonus fallax</i> (3)	“ <i>Bothrimonus sturionis</i> ” (2)	“ <i>Bothrimonus nylandicus</i> ” (4)
TL(mm)*	33.9 ± 18 (10.4–99.2)	51.8 ± 26.7 (16.2–103.6)	(5–20)	(5–130)	(20–170)	(8–90)	> 20
MW*	792 ± 370 (208–1800)	642 ± 238 (217–1042)	(800–1000)	(300–3000)	(1500–4500)	(400–2000)	(690–710)
TLsclx	513 ± 120 (279–737)	507 ± 134 (357–789)		(480–970)	1500	(135–680)	
MWsclx	616 ± 149 (389–853)	623 ± 266 (463–1147)		(570–2000)	(1400–1500)	(310–800)	
MTsclx	608 ± 165 (155–863)	605 ± 176 (421–863)		(560–1550)	(2000–2250)	(390–1140)	
DistPore*	500 ± 246 (205–1179)	781 ± 337 (253–1474)					
DistCirr*	501 ± 254 (188–1263)	735 ± 356 (212–1474)					
MDCirr	111 ± 28 (47–158)	98 ± 27 (48–147)		(125–230)	300	(120–200)	
Nprogs	78 ± 38 (27–183)	75 ± 25 (42–130)	(15–30)	(32–216)	(400–500)	(25–168)	(40–65)
Ntestes	28 ± 6 (12–40)	28 ± 8 (20–42)					
MDtestes	93 ± 20 (53–128)	87 ± 15 (63–116)		(70–110)	(100–190)	(45–75)	24
MLegg*	34 ± 4 (25–42)	37 ± 4 (31–44)	40	(33–46)	(35–41)	(33–42)	(30–34)
MWegg*	18 ± 2 (14–22)	20 ± 2 (16–23)	(25–30)	(20–29)	(27–31)	(21–30)	20
MLut	383 ± 167 (115–919)	440 ± 212 (126–836)					
MWut*	473 ± 182 (175–916)	379 ± 130 (140–533)					
MLov	312 ± 132 (81–660)	308 ± 97 (203–470)					
MWov*	451 ± 166 (191–912)	316 ± 122 (134–509)					
MWvt*	158 ± 78 (53–391)	111 ± 43 (73–239)					
Prout (%)	55 ± 12 (33–84)	56 ± 10 (36–70)					
Proov (%)	51 ± 14 (32–83)	49 ± 14 (32–83)					
Provt (%)	18 ± 8 (7–43)	19 ± 11 (9–49)		72	75	(46–55)	

molecular analyses, showed considerable overlap in the ranges of most characters, although the central genotype was generally greater in length and more slender on average. Indeed, significant differences ($P < 0.05$) were found between the total length (TL),

maximum width (MW), genital pore distances (PoreDist, CirrDist), egg mean width and length (MWegg, MLegg) and uterus, ovary and vitellarium mean width (MWut, MWov, MWvt, respectively) (Table 2). Morphometric comparisons also showed




Fig. 3. Scanning electron micrographs of the scolex of *Didymobothrium rudolphii* collected from *Solea lascaris* along the Portuguese coast. (A) Apical view of the scolex showing the longitudinal opening. (B) Scanning electron micrograph of a longitudinal section of the scolex showing the median septum. (C) Microtriches in longitudinal section, covering the body cut in transverse section. (D) Microtriches covering the surface of the scolex. BC, bothridial cavity; MS, median septum; MT, microtriches.

that most measurements and counts taken from both genotypes of *D. rudolphii* were within the range of those found for other Acrobothriidae (*sensu* Gibson, 1994), with the exception of those of *Bothrimonus fallax* Lühe, 1900 which were generally greater (Table 2). Minimum values in *D. rudolphii* (*sensu lato*) were generally lower than for those of *Diplocotyle olrikii*, but higher than those of '*Bothrimonus sturionis*' Duvernoy, 1842 (actually a species of *Diplocotyle* (see Gibson, 1994)), particularly those of the scolex (Table 2). Although few records were found for *D. nylandica*, the measurements of this species differed most from those of *Didymobothrium rudolphii*. The range of egg measurements rarely overlapped with those of the other species, with the eggs of *D. rudolphii* being typically smaller (Table 2).

Morphology and ultrastructure of Didymobothrium rudolphii 'common' genotype

Most features described are the same for the 2 genotypes of *D. rudolphii*. However, where given, the measurements describe only the 'common' genotype to avoid conflation of the 2 cryptic entities (see Table 2 for the morphometrics of the 'central' genotype). Similarly, all histological and SEM studies were conducted on specimens representing the 'common' genotype as present in the northern region.

Body elongate, dorso-ventrally flattened, 10–99 mm in length, 0.2–1.8 mm in width, with ~78 proglottides, most exhibiting the same degree of maturation. Scolex slightly rounded (Fig. 3A), comprising 2 strongly muscular, forwardly directed, hollow bothridia separated by complete septum




Fig. 4. Scanning electron micrographs and histological sections of the strobila of *Didymobothrium rudolphii* collected from *Solea lascaris* along the Portuguese coast. (A) Ventral view of the strobila. (B) Lateral view of the strobila showing the irregularly alternating pattern of genital pores. (C and D) Sagittal sections showing the reproductive organs and their openings. (E) Longitudinal section showing the cirrus-sac, vagina and uterus. CA, common atrium; CR, cirrus; CS, cirrus-sac; FP, female pore; UT, uterus; VG, vagina. Arrow indicates the anterior end of the worm.

(Fig. 3B). (Live worms presented significant variation in scolex shape, appearing to have either 1 or 2 hollow bothridia due to the contraction or extension of the septum (see also Burt and Sandeman, 1969)). Entire body, including scolex and internal surface of bothridia, covered with short, filiform microtriches $\sim 2 \mu\text{m}$ in length (Fig. 3C and D). Male and female genital pores open separately, in close proximity, along median line of strobila, most frequently on same side (Fig. 4A) but always irregularly alternating at some point in series (Fig. 4B). Male pore anterior to female pore; everted cirrus teat-like in shape (Fig. 4A and B), devoid of microtriches. Cirrus-sac ovoid (Fig. 4C); walls muscular (Fig. 4C and E). Female pore ovoid (Fig. 4A), with muscular common atrium into which vagina and uterus open side-by-side (Fig. 4D and E).

Proglottides exhibit same degree of maturation in most individuals (Figs 5A and 6A), but some worms presented less well-developed proglottides anteriorly (Figs 5B and 6B). Internal longitudinal musculature well developed, forming bundles between vitelline fields and medulla (Figs 5C and 6C). Vitellarium follicular and continuously distributed along margins of strobila (Figs 5D and 6A). Number of testes associated with each proglottis also varied (12–42) according to state of maturity. Vas deferens convoluted, often filled with spermatozoa, enters cirrus-sac forming ejaculatory duct (Fig. 5E). Vitellarium occupies 7–43% of body width depending on degree of maturation. Ovary medullary, lobulate (Fig. 5D), located posterior to uterus, occupies 32–83% of strobilar width, thin-walled, contains muscle fibres and glycogen (PAS-positive, diastase labile); when




Fig. 5. Light micrographs and histological sections of the reproductive organs of *Didymbothrium rudolphii*. (A) Mature proglottides, (B) immature proglottides, (C) sagittal section showing longitudinal muscle fibres and reproductive organs (uterus, cirrus-sac). (D) Transverse section showing vitelline follicles, testes and ovary, (E) male genital apparatus, and (F) mature ovary. CS, cirrus-sac; ED, ejaculatory duct; GP, genital primordial; LMB, longitudinal muscle bundles; OC, oocytes; OV, ovary; TT, testis; UT, uterus; VD, vas deferens; VT, vitelline follicles.




Fig. 6. Strobila of *Didymobothrium rudolphii*. (A) Mature region, (B) immature region, (C) longitudinal section showing muscle fibres and reproductive organs (uterus, ovary, cirrus-sac, testes). Arrow indicates the anterior end of the worm.

mature, all oocytes appear to be at similar stage of development (Fig. 5F).

Uterus tubular, convoluted (Fig. 7A), packed with eggs at varying degrees of maturity throughout its length (Fig. 7B–D); uterine field extends over 33–84% of strobilar width and overlaps ovary. Proximal uterus filled with oocytes surrounded by

thin, translucent shell, whereas eggs in middle and distal regions have glycogenic (PAS-positive, diastase labile) envelope (Fig. 7C); oblong eggs within most distal region of uterus have fully-developed tanned shells, including tuft of filaments at narrow pole (Fig. 7D–F) resembling ‘spaghetti’ (Fig. 7F) and containing glycogen (PAS-positive, diastase




Fig. 7. Scanning electron micrographs and histological sections of the uterus and eggs of *Didymobothrium rudolphii* collected from *Solea lascaris* along the Portuguese coast. (A) Sagittal section of the strobila showing the uterine shape, (B) section of the uterus filled with eggs, (C) different stages of oocytes and eggs along the ovary, (D) eggs surrounded by a mucous envelope and glycogen filaments, (E) egg outside attached to the individual's body, and (F) egg-shell. EG, egg; ES, egg-shell; FL, glycogen filaments; OC, oocytes; UT, uterus.

labile) which are expelled with the eggs (Fig. 7E). Eggs 25–42 μm in length, 14–22 μm in width; SEM revealed their rough surface (Fig. 7F).

Statistical analyses

According to the PCA ordination diagram (Fig. 8), some degree of morphological differentiation consistent with that of genetic differentiation was

found along the Portuguese coast, with most samples from the north and south (N and S) appearing on the lower half of the diagram and those from the centre (C) on the upper part of the diagram. Variables representing the major contribution to this pattern were the maximum width of the strobila (MW) and the distances between the genital pores (DistCirr, DistPore), although those associated with the scolex length, width and thickness (TLsclx, MWsclx and MTsclx, respectively), and those associated with the




Fig. 9. Seasonal variation in the mean intensity of *Didymobothrium rudolphii* (*sensu lato*) (bars) and the number of prey items found in the stomach contents of *Solea lascaris* for the 3 areas off the Portuguese coast sampled. Prey number was log transformed due to scaling constraints. Abbreviations: AmpN, number of Amphipoda; BivN, number of Bivalvia; CumN, number of Cumacea; DecN, number of Decapoda; DrudI, *D. rudolphii* mean intensity; MysN, number of Mysidacea; PolN, number of Polychaeta.

important components (Fig. 9). When the diet was analysed according to season and locality by means of a multivariate correspondence analysis, using the number or weight of the different prey groups, the Mysidacea was the most important group in the diet in the central region during summer and autumn. On the other hand, during spring in the central region and in the northern and southern regions throughout the year, Amphipoda and Bivalvia were more important (A. M. Pinheiro, personal communication).

DISCUSSION

Molecular sequence analyses readily differentiated 2 distinct entities of *Didymobothrium*, corroborating the earlier findings of Renaud and Gabrion (1988) based on allozyme data. Although no 'molecular yardstick' has yet been calculated to delineate taxonomic entities of different rank in the Cestoda (and may, in any case, prove to be of limited validity in isolation of other data), the degree of sequence divergence between the 'common' and 'central'

genotypes is considerable and easily in the range of that predicted for different species of the same genus (i.e. 1.9–2.1%). Within the Digenea, for example, ITS sequence divergences of 2–3% have been shown to correlate well with morphological differences used to circumscribe congeners by traditional means (Nolan and Cribb, 2005).

Generally, the external morphology of *D. rudolphii* described here is in agreement with that reported in the original description of this species by Monticelli (1890) and its subsequent redescrptions (Monticelli, 1892; Nybelin, 1922), especially with regard to the morphology of the bothridea and everted cirrus, the lack of external segmentation and the irregularly alternating genital pores. The present work provides the first description of aspects of the internal anatomy and egg shape of *D. rudolphii*, thus contributing to a better understanding of its likely life-cycle. For example, the finding of polar filaments on the eggs, together with its rough surface, suggests that they may have the facility of attachment, thus assisting transfer to the intermediate host.

The comparison of the morphological data for *D. rudolphii* with those from the better studied *Diplocotyle nylandica* and *D. olrikii* (see MacKinnon and Burt, 1984; Brunanska *et al.* 2005; Poddubnaya *et al.* 2005) and those from *Bothrimonus* spp., in conjunction with the genetic data, has shown that *Didymobothrium* can be differentiated from the other spathebothriideans, thus at least partly corroborating the systematic classification presented by Gibson (1994).

Although we cannot rule out sampling error given the small number of samples sequenced (i.e. 2 per season per area), the lack of co-occurrence of the 2 genotypes during the same season (despite being sympatric with regard to the central locality) is in stark contrast to the findings of Renaud and Gabrion (1988), who found the 2 forms to co-occur. They suggested that the sympatric existence of 2 species could be explained by differences in intermediate (presumably amphipod) host usage. Whereas one species of *Didymobothrium* (which they referred to as *Bothrimonus*) utilized an intermediate host species present throughout the year, the other utilized another, only present in spring and summer, overwintering in the egg stage. This life-cycle is similar to that described by Sandeman and Burt (1972) for '*B. sturionis*' (actually *Diplocotyle* (see Gibson, 1994)). Although such ecological niche separation could account for the divergence, it does not in itself explain why the 'common' form is absent for most of the year in the central region, while being present throughout the year in the north and south. This pattern requires one of a number of different potential isolating mechanisms: a method of active exclusion of the 'common' form by the 'central' form in the definitive host; a seasonal and ecological exclusion or exchange of intermediate host species

occurring in the central region only; or an extreme feeding preference of *Solea lascaris* for a particular species of intermediate host when available in the environment. As for most of the Soleidae, *S. lascaris* feeds on a wide range of small invertebrates, their importance in the diet being defined by their abundance in the environment (Link *et al.* 2005). The composition and seasonal variation of *S. lascaris* diet along the Portuguese coast found in this study, supported by findings of other investigations (Cabral *et al.* 2002; A. M. Pinheiro, unpublished data), revealed a preference for feeding on Mysidacea during summer and autumn in the central region, whereas Amphipoda were preferred in all other regions and seasons sampled. The finding of a change in diet concordant with the occurrence of *D. rudolphii* specimens presenting different genotypes suggests that Mysidacea could act as the intermediate host of the 'central' form of *D. rudolphii*. Moreover, mean intensity values during these seasons were the highest in the central region, whereas the lowest value was recorded when the consumption of Amphipoda was much higher. Thus, some ecological host switching might have occurred in the life-cycle of *D. rudolphii* in the central region off the Portuguese coast, either due to environmental pressure that might have diminished the availability of the original host (Amphipoda), or to the ecological pressure exerted by the feeding preferences of *S. lascaris*.

The existence of cryptic species of cestodes infecting Pleuronectiformes off the Portuguese coast may not be entirely unusual, as they have also been reported in the pseudophyllidean *Bothriocephalus scorpii* (Müller, 1776) (Renaud *et al.* 1986; Verneau *et al.* 1997). The 2 species of this complex, *B. renaudii* Ortega and Valero, 1989 and *B. gregarius* Renaud, Gabrion and Pasteur, 1983, parasitizing turbot, *Scophthalmus maximus*, have a disjunctive distribution – *B. renaudii* occurring in the Atlantic Ocean, including off the western Portuguese coast, and *B. gregarius* off the southern Portuguese coast and in the Mediterranean Sea, English Channel, Baltic Sea and North Sea (Renaud *et al.* 1986). Although the life-cycle of *Didymobothrium* is unlikely to be the same as that for *Bothriocephalus* spp., something in the behaviour of the definitive host may facilitate the formation of sibling species in cestodes that utilize these fish hosts.

Although the present data are sufficient to circumscribe 2 different species of *Didymobothrium* using molecular characters, a morphological diagnosis remains tenuous at best, and we are reluctant to make nomenclatural changes on the basis of DNA alone. However, the results of the morphological analyses suggest that species may be differentiated by their overall size, or length to width ratio, and that future studies guided by these results may show this to be a reliable character. More sampling needs to be

done in the central region in order to understand better how these entities remain sympatric but do not overlap in time or in the host infrapopulation, and to rule out the possibility that a sampling error has biased such conclusions. Elucidation of their life-cycles, presumed to involve amphipods and/or mysidaceans, could assist in explaining the speciation process.

This study was financed in part by the Fundação para a Ciência e a Tecnologia (FCT) through grants to J.F.M. (SFRH/BD/8983/2002) and M.J.S. (Grant SFRH/BSAB/492/2005), and by the Treaty of Windsor Programme 2005–06 (LIS/992/2). The authors thank M. Helena Sousa (Universidade do Porto, Portugal) for assistance with the histology, and A. Ball (NHM) and C.P. Santos (Instituto Oswaldo Cruz) for assistance with the SEM, including expert guidance to P.D.O. on freeze-fracture methods. P.D.O. thanks R. Kuchta and H. Brabec (University of South Bohemia, Czech Republic) and A. Shinn (University of Stirling, Scotland) for assistance with the collection of *Diplocotyle olrikii*, and T. Scholz (University of South Bohemia, Czech Republic) for providing ethanol-fixed samples of *Cyathocephalus*. The authors also thank C. Griffin (NHM) for assistance with sequencing problematic samples and E. Sherlock (NHM) for preparation of the molecular voucher specimens. Finally we thank the anonymous reviewers for their helpful comments.

REFERENCES

- Agustí, C., Aznar, F. J., Olson, P. D., Littlewood, D. T. J., Kostadinova, A. and Raga, J. A.** (2005). Morphological and molecular characterization of tetraphyllidean merocercooids (Platyhelminthes: Cestoda) of striped dolphins (*Stenella coeruleoalba*) from the western Mediterranean. *Parasitology* **130**, 461–474.
- Aznar, F. J., Agustí, C., Littlewood, D. T. J., Raga, J. A. and Olson, P. D.** (2007). The role of cetaceans in the tetraphyllidean life cycle: molecular and ecological data from the western Mediterranean. *International Journal for Parasitology* **37**, 243–255.
- Brickle, P., Olson, P. D., Littlewood, D. T. J., Bishop, A. and Arkhipkin, A.** (2001). Parasites of *Loligo gahi* from waters off the Falkland Islands with a molecular-based identification of their cestode larvae. *Canadian Journal of Zoology* **79**, 2289–2296.
- Brunanska, M., Poddubnaya, L. G. and Dezfali, B. S.** (2005). Vitellogenesis in two spathebothriidean cestodes. *Parasitology Research* **96**, 390–397.
- Burt, M. D. B. and Sandeman, I. M.** (1969). Biology of *Bothrimonus* (= *Diplocotyle*) (Pseudophyllidea: Cestoda) Part I. History, description, synonymy, and systematics. *Journal of the Fisheries Research Board of Canada* **26**, 975–996.
- Cabral, H. N., Lopes, M. and Loeper, R.** (2002). Trophic niche overlap between flatfishes in a nursery area in the Portuguese coast. *Scientia Marina* **66**, 293–300.
- Davydov, V. G., Poddubnaya, L. G. and Kuperman, B. I.** (1997). An ultrastructure of some systems of the *Diplocotyle olrikii* (Cestoda: Cyathocephalata) in relation to peculiarities of its life cycle. *Parazitologiya* **31**, 132–141. (In Russian.)
- Georgiev, B., Biserkov, V. and Genov, T.** (1986). *In toto* staining method for cestodes with iron acetocarmine. *Helminthologia* **23**, 279–281.
- Gibson, D. I.** (1994). Order Spathebothriidea Wardle and McLeod, 1952. In *Keys to the Cestode Parasites of Vertebrates* (ed. Khalil, L. F., Jones, A. and Bray, R. A.), pp. 15–19. CAB International, Wallingford.
- Gibson, D. I. and Valtonen, E. T.** (1983). Two interesting records of tapeworms from Finnish fishes. *Aquilo, Ser. Zoologica* **22**, 45–49.
- Hanzelová, V., Kuchta, R., Scholz, T. and Shinn, A. P.** (2005). Morphometric analysis of four species of *Eubothrium* (Cestoda: Pseudophyllidea) parasites of salmonid fish: an interspecific and intraspecific comparison. *Parasitology International* **54**, 207–214.
- Infante, C., Catanese, G. and Machado, M.** (2004). Phylogenetic relationships among ten sole species (Soleidae, Pleuronectiformes) from the Gulf of Cádiz (Spain) based on mitochondrial DNA sequences. *Marine Biotechnology* **6**, 612–624.
- Link, J. S., Fogarty, M. J. and Langton, R. W.** (2005). The trophic ecology of flatfishes. In *Flatfishes Biology and Exploitation* (ed. Gibson, R. N.), pp. 185–212. Blackwell Publishing, Oxford.
- Mackiewicz, J. S.** (2003). Caryophyllidea (Cestoidea): molecules, morphology and evolution. *Acta Parasitologica* **48**, 143–154.
- MacKinnon, B. M. and Burt, M. D. B.** (1984). The comparative ultrastructure of spermatozoa from *Bothrimonus sturionis* Duv. 1842 (Pseudophyllidae) *Pseudanthobothrium hansenii* Baer, 1956 (Tetraphyllidae), and *Monoecocestus americanus* Stiles, 1895 (Cyclophyllidea). *Canadian Journal of Zoology* **62**, 1059–1066.
- Maddison, W. P. and Maddison, D. R.** (2005). *MacClade: Analysis of Phylogeny and Character Evolution*. Sinauer Associates, Sunderland, Massachusetts.
- McGinnis, S. and Madden, T. L.** (2004). BLAST: at the core of powerful and diverse set of sequence analysis tools. *Nucleic Acids Research* **32**, W20–W25.
- Monticelli, S.** (1890). Note helminthologique. *Bollettino Società Naturalisti di Napoli* **4**, 189–208.
- Monticelli, S.** (1892). Sul genere *Bothrimonus*, Duvernoy e proposte per una classificazione dei Cestodi. *Monitore Zoologico Italiano* **5**, 100–108.
- Nolan, M. and Cribb, T. H.** (2005). The use and implications of ribosomal DNA sequencing for the discrimination of digenean species. *Advances in Parasitology* **60**, 102–156.
- Nybelin, O. N.** (1922). Anatomisch-systematische Studien über Pseudophyllideen. *Göteborgs Kungl. Vetenskaps-och Vitterhets-Samhälles Handlingar, Fjärde-Följden* **26**, 1–228.
- Nylander, J. A. A.** (2004). *MrModelTest, program distributed by the author*. Evolutionary Biology, Uppsala University, Sweden.
- Okaka, C. E.** (2000). Maturity of the procercoid of *Cyathocephalus truncatus* (Eucestoda: Spathebothriidae) in *Gammarus pulex* (Crustacea: Amphipoda) and the tapeworms life cycle using the amphipod as the sole host. *Helminthologia* **37**, 153–157.
- Olson, P. D., Cribb, T. H., Tkach, V. V., Bray, R. A. and Littlewood, D. T. J.** (2003). Phylogeny and

- classification of the Digenea (Platyhelminthes: Trematoda). *International Journal for Parasitology* **33**, 733–755.
- Olson, P. D., Littlewood, D. T. J., Griffiths, D., Kennedy, C. R. and Arme, C.** (2002). Evidence for the co-existence of separate strains or species of *Ligula* in Lough Neagh, Northern Ireland. *Journal of Helminthology* **76**, 171–174.
- Olson, P. D., Littlewood, D. T. J., Bray, R. A. and Mariaux, J.** (2001). Interrelationships and evolution of the tapeworms (Platyhelminthes: Cestoda). *Molecular Phylogenetics and Evolution* **19**, 443–467.
- Olson, P. D. and Tkach, V. V.** (2005). Advances and trends in the molecular systematics of the parasitic Platyhelminthes. *Advances in Parasitology* **60**, 165–243.
- Pertierra, A. A. G.** (2002). Redescription of *Proteocephalus bagri* and *P. rhamdiae* (Cestoda: Proteocephalidae), parasites of *Ramdia quelen* (Siluriformes: Pimelodidae) from South America, with comments on morphological variation. *Folia Parasitologica* **49**, 55–66.
- Poddubnaya, L. G., Mackiewicz, J. S. and Kuperman, B. I.** (2003). Ultrastructure of *Archigetes sieboldi* (Cestoda: Caryophyllidea): relationship between progenesis, development and evolution. *Folia Parasitologica* **50**, 275–292.
- Poddubnaya, L. G., Mackiewicz, J. S., Brunanska, M. and Scholz, T.** (2005). Ultrastructural studies on the reproductive system of progenetic *Diplocotyle olrikii* (Cestoda, Spathebothriidae): Ovarian tissue. *Acta Parasitologica* **50**, 199–207.
- Poddubnaya, L. G., Gibson, D. I., Swiderski, Z. and Olson, P. D.** (2006). Vitellocyte ultrastructure in the cestode *Didymobothrium rudolphii* (Monticelli, 1890): possible evidence for the recognition of divergent taxa within the Spathebothriidae. *Acta Parasitologica* **51**, 255–263.
- Posada, D. and Crandall, K. A.** (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**, 817–818.
- Protasova, E. N. and Roytman, V. A.** (1995). [Cyathocephalates, tapeworm helminths of marine and freshwater fish (Cestoda: Pseudophyllidea: Cyathocephalata).] *Osnovy Tsestodologii*, Vol. 12. Institute of Parasitology, Russian Academy of Sciences, Moscow.
- Renaud, F. and Gabrion, C.** (1988). Speciation in Cestoda: evidence for two sibling species in the complex *Bothrimonus nylandicus* (Schneider 1902) (Cestoda: Cyathocephalidae). *Parasitology* **97**, 139–147.
- Renaud, F., Gabrion, C. and Pasteur, N.** (1986). Geographical divergence in *Bothriocephalus* (Cestoda) of fishes demonstrated by enzyme electrophoresis. *International Journal for Parasitology* **16**, 553–558.
- Reyda, F. B. and Olson, P. D.** (2003). Cestodes of cestodes of Peruvian freshwater stingrays. *Journal of Parasitology* **89**, 1018–1024.
- Ronquist, F. and Huelsenbeck, J.** (2003). MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- Sandeman, I. M. and Burt, M. D. B.** (1972). Biology of *Bothrimonus* (= *Diplocotyle*) (Pseudophyllidea: Cestoda): ecology, life cycle, and evolution; a review and synthesis. *Journal of Fisheries Research Board of Canada* **29**, 1381–1395.
- Schneider, G.** (1902). *Bothrimonus nylandicus* n. sp. *Archiv für Naturgeschichte* **1**, 72–78.
- Swofford, D. L.** (2001). *PAUP*. Phylogenetic Analysis Using Parsimony (*and other Methods). Version 4*. Sinauer Associates, Massachusetts.
- SPSS 13.0** (2004). SPSS Inc., Chicago.
- ter Braak, C. J. F. and Smilauer, P.** (2002). *Canoco for Windows Version 4.5*. Biometris – Plant Research International, Wageningen, The Netherlands.
- Verneau, O., Renaud, F. and Catzeffis, F.** (1997). Evolutionary relationships of sibling tapeworm species (Cestoda) parasitizing teleost fishes. *Molecular Biology and Evolution* **14**, 630–636.
- Zehnder, M. P. and de Chambrier, A.** (2000). Morphological and molecular analyses of the genera *Peltidocotyle* Diesing, 1850 and *Othinoscotyle* Woodland, 1933, and morphological study of *Woodlandiella* Freze, 1965 (Eucestoda, Proteocephalidae), parasites of South American siluriform fishes (Pimelodidae). *Systematic Parasitology* **46**, 33–43.