

1. Hammond NE, Saxena MK, Taylor C, et al. Temperature management of non-elective intensive care patients without neurological abnormalities: a point prevalence study of practice in Australia and New Zealand. *Crit Care Resusc* 2013;15:228-33.
2. Saxena MK, Hammond NE, Taylor C, et al. A survey of fever management for febrile intensive care patients without neurological injury. *Crit Care Resusc* 2011;13:238-43.
3. Young P, Saxena M, Eastwood GM, Bellomo R, Beasley R. Fever and fever management among intensive care patients with

- known or suspected infection: a multicentre prospective cohort study. *Crit Care Resusc* 2011;13:97-102.
4. Niven DJ, Gaudet JE, Laupland KB, Mrklas KJ, Roberts DJ, Stelfox HT. Accuracy of peripheral thermometers for estimating temperature: a systematic review and meta-analysis. *Ann Intern Med* 2015;163:768-77.
 5. Jefferies S, Braithwaite I, Walker S, et al. Randomized controlled trial of the effect of regular paracetamol on influenza infection. *Respirology* 2016;21:370-7.
DOI: 10.1056/NEJMc1601361

Malignant Transformation of *Hymenolepis nana* in a Human Host

TO THE EDITOR: Muehlenbachs et al. (Nov. 5 issue)¹ describe a case of malignant transformation in a cestode, *Hymenolepis nana*, and transformation resulting in individual proliferating cells invading host tissues. The authors cite some earlier cases as examples of proliferating parasites that maintained recognizable metazoan tissues. Although their report is mostly accurate, it understates the century-long history of numerous reports of transformations among cestodes that infect humans and other animals, with typically fatal outcomes. Earlier reports used different terminology that is more consistent with invertebrate zoology, such as “aberrant” or “anomalous,” but the developmental and morphologic criteria for neoplasms and malignant transformations would apply as well. Most recent are our descriptions of aberrant interepithelial and epitheliomesenchymal interactions resulting in transformation into abnormal new (neoplastic) structures in metacestodes, including mesocestoides² and spirometra species,³ thus producing the malignant proliferative disseminated lethal infections previously reported in many host species.⁴ In retrospective analysis, the transformations we reported appear to be identical to all known cases, including the earliest human cases in Japan in 1905 and Florida in 1908.⁵

David B. Conn, Ph.D.

Harvard University Museum of Comparative Zoology
Cambridge, MA
bconn@berry.edu

No potential conflict of interest relevant to this letter was reported.

1. Muehlenbachs A, Bhatnagar J, Agudelo CA, et al. Malignant transformation of *Hymenolepis nana* in a human host. *N Engl J Med* 2015;373:1845-52.
2. Conn DB, Galán-Puchades MT, Fuentes MV. Interactions between anomalous excretory and tegumental epithelia in aberrant Mesocestoides tetrathyridia from *Apodemus sylvaticus* in Spain. *Parasitol Res* 2010;106:1109-15.
3. Nobrega-Lee M, Hubbard G, Loverde P, et al. Sparganosis in wild-caught baboons (*Papio cynocephalus anubis*). *J Med Primatol* 2007;36:47-54.

4. McFarlane D, Mann KA, Harmon BG, Williamson LH, Conn DB. Pleural effusion secondary to aberrant metacestode infection in a horse. *Compend Contin Educ Pract Vet* 1994;16:1032-5.
5. Stiles CW. The occurrence of a proliferating cestode larva (*Sparganum proliferum*) in man in Florida. *Hyg Lab Bull* 1908;40:7-18.
DOI: 10.1056/NEJMc1600490

THE AUTHORS REPLY: Conn notes the long history of atypical infections with tapeworm larvae. We agree that many of these aberrant forms probably share an underlying disease process. We urge caution in considering them as one entity, however, since malformed larvae are typically not considered to be neoplasms despite rapid cellular proliferation. Moreover, atypical infections often occur in atypical hosts or immunocompromised patients, as in the most similar previous case reports.¹ This fact suggests that the absence of normal host-defense signals stimulates abnormal tapeworm growth,² tissue dissemination, and ultimately neoplasia. We propose that these infections represent a spectrum of disease (as in cancer in humans), from the stimulated proliferation of normal tapeworm stem cells to the acquisition of somatic mutations, clonal overgrowth, and frank neoplasia (i.e., from low-grade to high-grade malignant transformation). Fundamental questions remain: Do atypical proliferative infections contain somatic mutations? Have true neoplasms been observed previously? Do somatic mutations contribute to aggressive behavior in more common tapeworm infections, including racemose cysticercosis and late-stage echinococcosis? Our finding that aberrant tapeworm development can include neoplasia lends credence to these inquiries.

Atis Muehlenbachs, M.D., Ph.D.

Blaine A. Mathison, B.S.M.(A.S.C.P.)

Centers for Disease Control and Prevention
Atlanta, GA
vkd6@cdc.gov

Peter D. Olson, Ph.D.

Natural History Museum
London, United Kingdom

The findings and conclusions in this letter are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Since publication of their article, the authors report no further potential conflict of interest.

1. Olson PD, Yoder K, Fajardo L-G LF, et al. Lethal invasive cestodiasis in immunosuppressed patients. *J Infect Dis* 2003; 187:1962-6.

2. Ito A. Basic and applied problems in developmental biology and immunobiology of cestode infections: *Hymenolepis*, *Taenia* and *Echinococcus*. *Parasite Immunol* 2015;37:53-69.

DOI: 10.1056/NEJMc1600490

Thyroid Nodules

TO THE EDITOR: Burman and Wartofsky (Dec. 10 issue)¹ describe a clinical approach to thyroid nodules and indicate that a “radioisotopic (iodine-123 or technetium-99m) scan with measurement of radioisotope uptake to confirm autonomous function is indicated only if the thyrotropin level is suppressed.” We think that a thyroid radioisotope scan may be considered in patients with findings on thyroid fine-needle aspiration that are suggestive of a follicular neoplasm, particularly if the thyrotropin level is in the lower end of the normal range. Although there is a low likelihood of finding a functioning nodule when thyrotropin levels are not suppressed, this approach might spare some patients from undergoing unnecessary surgery.

Our suggestion is consistent with the revised 2009 guidelines of the American Thyroid Association (ATA) on thyroid nodules.² The 2015 ATA guidelines, however, remained silent about this issue.³

Abdallah M. Beano, M.D.

Mohammad A. Zmaili

Ayman A. Zayed, M.D.

University of Jordan School of Medicine
Amman, Jordan
baraayman@gmail.com

No potential conflict of interest relevant to this letter was reported.

1. Burman KD, Wartofsky L. Thyroid nodules. *N Engl J Med* 2015;373:2347-56.

2. American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2009;19:1167-214.

3. Haugen BRM, Alexander EK, Bible KC, et al. 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2016;26:1-133.

DOI: 10.1056/NEJMc1600493

TO THE EDITOR: In their review of palpable thyroid nodules, Burman and Wartofsky do not discuss incidentally detected nodules and only

briefly address overdiagnosis and overtreatment of incidentally detected papillary thyroid cancers. These lesions outnumber palpable cancers by orders of magnitude in clinical practice.¹

The current guideline of the American College of Radiology was not mentioned in the article. It calls for no follow-up at all for incidentally detected nodules that are smaller than 1.0 cm in diameter in patients who are younger than 35 years of age or smaller than 1.5 cm in diameter in patients who are 35 years of age or older.²

Readers and patients would benefit from awareness of the low probability of cancer in small, incidentally detected nodules. They also would benefit from knowing that with appropriate evaluation and therapy, a favorable prognosis is associated with nodules that are found to be histologically malignant.

David Seidenwurm, M.D.

Jonathan Breslau, M.D.

Sutter Medical Group
Sacramento, CA
breslaj@sutterhealth.org

No potential conflict of interest relevant to this letter was reported.

1. Tan GH, Gharib H. Thyroid incidentalomas: management approaches to nonpalpable nodules discovered incidentally on thyroid imaging. *Ann Intern Med* 1997;126:226-31.

2. Hoang JK, Langer JE, Middleton WD, et al. Managing incidental thyroid nodules detected on imaging: white paper of the ACR Incidental Thyroid Findings Committee. *J Am Coll Radiol* 2015;12:143-50.

DOI: 10.1056/NEJMc1600493

THE AUTHORS REPLY: The combined guidelines of the American Association of Clinical Endocrinologists–Associazione Medici Endocrinologi–European Thyroid Association do not recommend the performance of a radioisotope scan when findings on thyroid fine-needle aspiration are suggestive of a follicular neoplasm.¹ The 2015 ATA guidelines have removed this recommendation,² which was present in the 2009 ATA guidelines and based on expert opinion.