

Molecules and morphology reveal cryptic variation among digeneans infecting sympatric mullets in the Mediterranean

I. BLASCO-COSTA^{1*}, J. A. BALBUENA¹, J. A. RAGA¹, A. KOSTADINOVA^{2,3}
and P. D. OLSON⁴

¹Marine Zoology Unit, Cavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, PO Box 22 085, 46071 Valencia, Spain

²Institute of Parasitology, Biology Centre, Academy of Sciences of the Czech Republic, Branišovská 31, 370 05 České Budějovice, Czech Republic

³Central Laboratory of General Ecology, Bulgarian Academy of Sciences, 2 Gagarin Street, 1113 Sofia, Bulgaria

⁴Department of Zoology, Natural History Museum, Cromwell Road, London SW7 5BD, UK

(Received 13 July 2009; revised 8 August 2009; accepted 8 August 2009; first published online 23 October 2009)

SUMMARY

We applied a combined molecular and morphological approach to resolve the taxonomic status of *Saccocoelium* spp. parasitizing sympatric mullets (Mugilidae) in the Mediterranean. Eight morphotypes of *Saccocoelium* were distinguished by means of multivariate statistical analyses: 2 of *Saccocoelium obesum* ex *Liza* spp.; 4 of *S. tensum* ex *Liza* spp.; and 2 (*S. cephalis* and *Saccocoelium* sp.) ex *Mugil cephalus*. Sequences of the 28S and ITS2 rRNA gene regions were obtained for a total of 21 isolates of these morphotypes. Combining sequence data analysis with a detailed morphological and multivariate morphometric study of the specimens allowed the demonstration of cryptic diversity thus rejecting the hypothesis of a single species of *Saccocoelium* infecting sympatric mullets in the Mediterranean. Comparative sequence analysis revealed 4 unique genotypes, thus corroborating the distinct species status of *Saccocoelium obesum*, *S. tensum* and *S. cephalis* and a new cryptic species ex *Liza aurata* and *L. saliens* recognized by its consistent morphological differentiation and genetic divergence. However, in spite of their sharp morphological difference the 2 morphotypes from *M. cephalus* showed no molecular differentiation and 4 morphotypes of *S. tensum* were genetically identical. This wide intraspecific morphological variation within *S. tensum* and *S. cephalis* suggests that delimiting species of *Saccocoelium* using solely morphological criteria will be misleading.

Key words: Digenea, Haploporidae, *Saccocoelium*, Mugilidae, cryptic species, molecules, morphology, rDNA.

INTRODUCTION

The adoption of molecular techniques in taxonomy has revealed unexpected genetic diversity within species throughout the tree of life (Bickford *et al.* 2007). This has been especially true for the Digenea, a large group of internal metazoan parasites of animals characterized by a substantial interspecific homogeneity of the morphological characters used for species discrimination. Although the discovery of cryptic species through molecular means does not require *a priori* knowledge of the morphological variation in the system under study (e.g. Donald *et al.* 2004; Miura *et al.* 2005; Leung *et al.* 2009), such knowledge is important for revealing characters that aid future identifications and thus promote a taxonomic change. Moreover, whereas many studies report the occurrence of unique genetic strains,

comparatively few include formal taxonomic revisions, as such changes require a significant understanding of the morphology and systematic history of the group in question. In the taxonomy of the digeneans, as elsewhere, there has been a clear trend toward combining molecular and morphological approaches (see Nolan and Cribb, 2005; Olson and Tkach, 2005) and this, along with increased sampling effort in recent years, has resulted in the discovery of cryptic species and higher taxa in at least 19 digenean families (e.g. Jousson and Bartoli, 2001; Nolan and Cribb, 2004, 2006; Chambers and Cribb, 2006; Miller and Cribb, 2007*a,b*; see Nolan and Cribb, 2005 for a recent review).

The Haploporinae Nicoll, 1914 (Digenea: Haploporidae) is a small group of poorly known haploporid digeneans which parasitize marine or brackishwater mugilid fishes (Mugilidae). Most species were described from the Mediterranean by Looss (1902) and this resulted in the erection of the bulk of the haploporine genera from this area. The most 'species-rich' of these genera, *Saccocoelium* Looss, 1902, was erected for 2 species, *S. obesum* Looss, 1902 (type-species) and *S. tensum* Looss, 1902,

* Corresponding author: Marine Zoology Unit, Cavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, PO Box 22 085, 46071 Valencia, Spain. Tel: +34 963543685. Fax: +34 963543733. E-mail: M.Isabel.Blasco@uv.es

discovered in the Adriatic Sea. Further studies have described and assigned 8 species to this genus: 6 nominal plus 2 transferred from *Lecithobotrys* Looss, 1902 (see Blasco-Costa *et al.* 2009*e* for details). However, most *Saccocoelium* spp. are only known from their original descriptions and, although the historically 'oldest' species described by Looss are the most widely reported in virtually all Mediterranean mullets, there are few records providing data on their morphology, and many authors have considered them synonymous (Dawes, 1947; Mikailov, 1958; Fischthal and Kuntz, 1963; Ferreti and Paggi, 1965; Moravec and Libosvářský, 1975).

Until recently, no attempts have been made to address the taxonomic diversity and morphological variability within the Haploporidae and this perhaps has led to the fact that a hypothesis for a single polymorphic species has been suggested in the case of Mediterranean representatives of 3 of the 5 genera of the Haploporinae (synonymies discussed in detail by Blasco-Costa *et al.* 2009*b, c, e*). In a series of taxonomic studies based on newly-collected material from mullets in the Mediterranean basin and a critical evaluation of the published data, we have re-assessed the status of the constituent species of the 'Mediterranean' genera of the Haploporinae erected by Looss (1902); these studies have resulted in the erection of 3 new genera and descriptions of 4 new species (Blasco-Costa *et al.* 2009*b, c, d*). The revision of *Saccocoelium*, in particular, led to the descriptions of 2 new species from *Mugil cephalus* L. (*Saccocoelium cephalis* Blasco-Costa *et al.*, 2009 and *S. currani* Blasco-Costa *et al.*, 2009), re-descriptions of *S. obesum* and *S. tensum* based on material from *Liza* spp. and the discrimination of the 4 species from the Mediterranean with the aid of multivariate analyses. We considered *S. portsaidensis* El-Shahawi, El-Gindy, Imam & Al-Bassel, 1992, *S. saoudi* El-Shahawi, El-Gindy, Imam & Al-Bassel, 1992 and *Neosaccocoelium aegyptiacus* El-Shahawi, El-Gindy, Imam & Al-Bassel, 1992 to be synonyms of *S. tensum*, thus indicating a wide intraspecific morphological variation within this species (Blasco-Costa *et al.* 2009*e*). Furthermore, in our samples from the western Mediterranean, we could distinguish morphologically 4 morphotypes of the latter species, 2 morphotypes of *S. obesum* and a third form, *Saccocoelium* sp., parasitizing *M. cephalus*, which, although not identified to the species level, differed morphologically from the species of *Saccocoelium* that we considered valid. These findings, in combination with the unexpected diversity of haploporids revealed in *M. cephalus* (see also Blasco-Costa *et al.* 2009*c*) prompted us to corroborate our findings using molecular data. Here we use morphological and DNA sequence data of the 28S and ITS2 rRNA genes to elucidate the taxonomic status of *Saccocoelium* spp. parasitizing mullets in the western Mediterranean. In particular, we test for the presence of

cryptic species *via* the assessment of a series of isolates of the suspected morphotypes of *S. obesum*, *S. tensum* and *S. cephalis* and examine the phylogenetic affinities of the forms parasitizing *M. cephalus* and *Liza* spp. The system selected for study is small, but characteristic for the known size of the genera of the Haploporidae, and thus provides an example for the application of a combined approach to species diversity within this problematic digenean group.

MATERIALS AND METHODS

Morphological data

Four sympatric mullet species, *Mugil cephalus*, *Liza aurata* (Risso), *L. ramado* (Risso) and *L. saliens* (Risso), were collected at 3 localities along the Mediterranean coast of Spain: Ebro Delta (40°30'–40°50'N, 0°30'–1°10'E); off Santa Pola (38°00'–38°20'N, 0°10'–0°40'E); and in a brackishwater lagoon near Santa Pola. Trematodes were killed in near-boiling saline solution, fixed in 70% alcohol, stained with iron acetocarmine, dehydrated through a graded alcohol series, cleared in dimethyl phthalate and examined as permanent mounts in Canada balsam. Eight morphotypes of *Saccocoelium* were distinguished on morphological grounds and sequenced in the course of the study: 2 of *S. obesum* *ex Liza* spp.; 4 of *S. tensum* *ex Liza* spp.; and 2 *ex M. cephalus* (*S. cephalis* and *Saccocoelium* sp.). The type- and voucher material is deposited in the British Museum (Natural History) Collection at the Natural History Museum, London (BMNH) (see Table 1 for Accession numbers). All measurements are in micrometres.

Statistical analyses

Multivariate statistical analyses were performed on 13 metrical variables. First, a principal components analysis (PCA) was applied to reveal the multivariate relationship between the specimens (excluding 'morphotype 4' of *S. tensum* represented by a single specimen). Secondly, a linear discriminant analysis (LDA; backward stepwise procedure) was applied to 51 specimens assigned to 7 *a priori* groups in order to evaluate the morphometric differences between them and to identify the variables yielding optimal separation. The squared Mahalanobis distances between the group centroids obtained in the LDA were then used to perform a test for association between genetic and morphological distances separating recognized morphotypes by applying a method based on the permutation of distance matrices. A Mantel test for matrix correlation was carried out by regressing the Mahalanobis distances in the distance matrix from the LDA on the distances in the percentage of sequence difference matrix (ITS2 region only). The

Table 1. *Saccocoelium* spp. and ougroup taxa sequenced, their hosts, localities, number of sequenced isolates for the two rDNA regions and EMBL/GenBank and BMNH Accession numbers for sequences and morphological vouchers, respectively

Species	Host	Locality	No. of isolates and EMBL/GenBank Accession numbers			
			28S	ITS2	BMNH Accession numbers	
<i>Saccocoelium brayi</i> n. sp.	<i>Liza saliens</i>	Ebro Delta	1	FJ211234	FJ211244	2008.10.7.77 (holotype); 2008.10.7.78-83 (paratypes)
<i>Saccocoelium cephalii</i>	<i>Mugil cephalus</i>	Ebro Delta	1	FJ211233	FJ211243	2008.10.7.23 (holotype); 2008.10.7.2-25 (paratypes)
<i>Saccocoelium obesum</i>	<i>Liza aurata</i> / <i>Liza ramado</i>	Ebro Delta	3/1	FJ211260/ FJ211159	FJ211266/ FJ211265	2008.10.7.38-39 (large morph 1)
<i>Saccocoelium tensum</i>	<i>Liza aurata</i> / <i>Liza ramado</i>	Ebro Delta Santa Pola (lagoon) Santa Pola (sea)	1/6	FJ211258/ FJ211257	FJ211264/ FJ211263	2008.10.7.41-44 (morphotype 1) 2008.10.7.45-46 (morphotype 2) 2008.10.7.47 (morphotype 3) 2008.10.7.48 (morphotype 4)
<i>Saccocoelium</i> sp.	<i>Mugil cephalus</i>	Ebro Delta	8	GQ357169	GQ357170	2008.10.7.49-51
<i>Forticulcita gibsoni</i>	<i>Mugil cephalus</i>	Santa Pola (sea)	1	FJ211239	FJ211249	2008.10.7.61 (holotype) 2008.10.7.62-76 (paratypes)
<i>Haploporus benedeni</i>	<i>Liza ramado</i>	Santa Pola (sea)	4	FJ211237	FJ211247	2008.10.7.52-55

significance of the best regression model was tested with a randomization approach (9999 random permutations of the dependent variable matrix) (Manly, 1997) using RT 2.1 program (Western EcoSystems Technology, Inc., Cheyenne, Wyoming).

Molecular data

Specimens were fixed live in 100% ethanol and stored at -20°C . Specimens were subsequently transferred into 300 μl of TNES urea (10 mM Tris-HCl (pH 8), 125 mM NaCl, 10 mM EDTA, 0.5% SDS, 4 M urea). Genomic DNA (gDNA) was extracted from single specimens using a phenol-chloroform protocol as described by Holzer *et al.* (2004). Alternatively, 1 M Tris-EDTA (pH 8) buffer was used to replace ethanol from the tissue of some specimens and gDNA was extracted using Qiagen® DNeasy™ tissue kit following the manufacturer's protocol, except for the proteinase-K incubation period, which was extended overnight, and the gDNA was further concentrated to a volume of $\sim 30 \mu\text{l}$ using Millipore Microcon® columns. Complete ITS2 rDNA sequences were amplified using primers 3S (5'-GTA CCG GTG GAT CAC GTG GCT AGT G-3'; Anderson and Barker, 1993) and ITS2.2 (5'-CCT GGT TAG TTT CTT TTC CTC CGC-3'; Anderson and Barker, 1993). Partial (domains D1–D3; ~ 1400 bps) 28S rDNA sequences were amplified using primers LSU5 (5'-TAG GTC GAC CCG CTG AAY TTA AGC A-3'; Littlewood *et al.* 2000) and LSU1500R (5'-GCT ATC CTG AGG GAA ACT TCG-3'; Tkach *et al.* 1999) or (~ 1600 bps) U178 (5'-GCA CCC GCT GAA YTT

AAG-3'; Lockyer *et al.* 2003) and L1642 (5'-CCA GCG CCA TCC ATT TTC A-3'; Lockyer *et al.* 2003). Polymerase chain reaction (PCR) amplifications were carried out using Ready-To-Go™ (Amersham Pharmacia Biotech) PCR beads (each containing ~ 1.5 units *Taq* DNA polymerase, 10 mM Tris-HCl (pH 9), 50 mM KCl, 1.5 mM MgCl_2 , 200 μM of each dNTP and stabilizers, including BSA), 20–70 ng of template DNA and 10 mM of each PCR primer. The following thermocycling profile was used for ITS2 rDNA amplification: denaturation of DNA (95°C for 3 min); 35 cycles of amplification (94°C for 50 sec, 54°C for 50 sec and 72°C for 1 min 20 sec); and 4 min extension hold at 72°C . The same profile but with annealing temperatures of 58°C and 56°C , respectively for the primer combinations LSU5-L1500R and U178-L1642 was applied for 28S rDNA amplification. PCR amplicons were either gel-excised or purified directly using Qiagen QIAquick™ PCR Purification Kit and cycle-sequenced from both strands using ABI BigDye™ Terminator v3.1 Ready Sequencing Kit, alcohol-precipitated, and run on an ABI 3730 automated sequencer. The PCR primers and internal primers 300F (5'-CAA GTA CCG TGA GGG AAA GTT G-3'), ECD2 (5'-CTT GGT CCG TGT TTC AAG ACG GG-3') and LSU1200R (5'-GCA TAG TTC ACC ATC TTT CGG-3') (Littlewood *et al.* 2000) in the case of the 28S rDNA products, were used for cycle sequencing. Contiguous sequences were assembled and edited using either Bioedit v7.0.5. (©1997–2005, Hall, 1999) or Sequencher™ (GeneCodes Corp., ver. 3.1.1) and submitted to GenBank (see Table 1 for Accession numbers).

Alignment and phylogenetic analysis

Newly generated ITS2 rDNA and partial 28S rDNA sequences of *Saccocoelium* spp. were aligned using MUSCLE (Edgar, 2004) together with the new sequences of the haploporines *Haploporus benedeni* Looss, 1902 and *Forticulcita gibsoni* Blasco-Costa *et al.*, 2009 (used as outgroup taxa) and adjustments made by eye using MacClade 4.08 (Maddison and Maddison, 2005). Sequences for both gene fragments were concatenated and regions of ambiguous alignment were defined in a character exclusion set. Pairwise distances for each rDNA region were calculated from the trimmed (to match the shortest sequence) aligned sequences with the absolute pairwise character difference (gaps treated as missing data) and the percentage of pairwise character differences on a total of 451 and 1189 unambiguously aligned positions for the ITS2 and the 28S rDNA, respectively.

ITS2 and 28S rDNA data partitions were analysed individually and combined using the methods of maximum parsimony (MP) and Bayesian inference (BI). MP analyses were performed with PAUP* 4.0b10 (Swofford, 2002) using a heuristic search strategy with 1000 search replicates, random-addition taxa sampling, tree-bisection-reconnection branch-swapping, with all characters run unordered with equal weights and with gaps treated as missing data. Nodal support was estimated by bootstrap analysis (heuristic search strategy with 1000 pseudo-replicates and 100 random sequence addition each). BI analyses were conducted using MrBayes version 3.1.2 (Ronquist and Huelsenbeck, 2003). Prior to the analyses, the nucleotide substitution model was estimated using ModelTest 3.06 (Posada and Crandall, 1998) independently for each data partition. The model GTR+ Γ (general-time-reversible model including gamma distributed among-site rate variation) was estimated as the one fitting the data best. The analyses were run over 1 million generations with a sampling frequency of 100. Consensus trees with mean branch lengths were constructed after log-likelihood values and substitution parameters plateaued at approximately generation number 12700 and 6600 for the ITS2 and 28S rDNA regions, respectively; and at 6400 for the combined analysis. Nodal support was estimated as posterior probabilities (Huelsenbeck *et al.* 2001).

RESULTS

Morphological data

Two morphotypes of S. obesum ex *Liza* spp. A detailed description of *S. obesum* based on material of 2 morphotypes (labelled 'large morph 1' and 'large morph 2' from *L. aurata* in the Ebro Delta and the Black Sea, respectively) is provided in Blasco-Costa *et al.* (2009e). In the latter morphological study, a third morphotype (labelled 'small morph') was

distinguished in *L. aurata* from the Ebro Delta, which indicated a possible existence of a cryptic species, although no new species was proposed. Examination and sequencing of additional material from *L. saliens* from the same locality confirmed the distinctness of the 'small morph', for which we propose the name *S. brayi* n. sp.

S. brayi n. sp.

Syn. *Saccocoelium* n. sp. of Blasco-Costa *et al.* (2009a)

Type-host: *Liza aurata* (Risso), the golden grey mullet (Mugilidae).

Other host: *Liza saliens* (Risso).

Type-locality: Off the Mediterranean coast of Spain, Ebro Delta (40°30'–40°50'N, 0°30'–1°10'E).

Site: Pyloric caeca of intestine.

Type-material: Holotype BMNH 2008.10.7.77; paratypes BMNH 2008.10.7.78–83.

Representative DNA sequences: GenBank nos. FJ211234 (partial D1–D3 28S rDNA) and FJ211244 (ITS2 rDNA); these were referred to as *Saccocoelium* n. sp. by Blasco-Costa *et al.* (2009a) to avoid nomenclatural problems due to uncertainty concerning the first publication of the name.

Etymology: The species is named for Dr Rodney A. Bray of the Natural History Museum, London, in recognition of his contributions to platyhelminth taxonomy, systematics and evolution.

Description (Fig. 1A–C)

(Based on 10 whole-mounted adult specimens; measurements in Table 2). Body elongate-oval, plump, with bell-shaped concavity at posterior extremity, with bluntly rounded posterior extremity and maximum width at posterior level of ventral sucker; width 43–59% of body length. Pre-oral lobe strongly developed, 25–51. Tegument thick (5–8), armed with large (8–10) sharp spines reaching to posterior extremity. Eye-spot pigment abundant, dispersed on either side of anterior part of pharynx and oral sucker. Oral sucker spherical, with ventral aperture. Ventral sucker cup-shaped, similar in size to or slightly larger than oral sucker (sucker length ratio 1:0.87–1.09; width ratio 1:1.00–1.28), in second quarter of body. Forebody relatively short, 35–41% of body length. Pre-pharynx absent or short (PL/PHL=0–0.6); pharynx strongly muscular, elongate-oval, larger than oral sucker. Oesophagus similar in length to pharynx. Intestinal bifurcation at level of posterior margin of ventral sucker; caeca 2 sac-like, large, with thick lining of cells, end blindly in about middle of hindbody at 26–33% from posterior extremity.

Testis single, median to dextral, subglobular, smooth, in last quarter of body; post-testicular field




Fig. 1. *Saccocoelium brayi* n. sp. ex *Liza aurata* and *L. saliens*. (A) Holotype (slightly flattened) ex *L. aurata*, ventral view with uterus in outline. (B) Paratype ex *L. saliens*, ventral view with uterus in outline. (C) Paratype ex *L. aurata*, detail of the terminal genitalia. Scale bars: A and B = 200 μm ; C = 100 μm .

9–23% of body length. External seminal vesicle saccular, subglobular, thin-walled (<2), smaller than internal seminal vesicle. Hermaphroditic sac massive, thick-walled (5–6), muscular, elongate-oval, reaches up to distance equal to length of ventral sucker posteriorly into hindbody, much longer than ventral sucker (HSL/VSL = 189–227%), contains internal seminal vesicle, numerous small prostatic cells, metraterm and hermaphroditic duct. Hermaphroditic duct wide (c.51–89), faintly-muscular, thick-walled; walls lined with crescentic, sclerotized structures. Internal seminal vesicle lined by single layer of cells, saccular, elongate-oval, occupies up to half of hermaphroditic sac. Genital atrium with strongly developed muscular wall. Genital pore median, at posterior level of pharynx.

Ovary round to transverse-oval, median to dextral, separated (holotype) from or contiguous (paratypes) with hermaphroditic sac, posterior margin of ventral sucker and testis, overlapping latter dorsally in some specimens. Uterine seminal receptacle distinct; blind seminal receptacle absent. Mehlis' gland and Laurer's canal not observed. Uterus thin-walled, occupies almost entire hindbody; metraterm distinct, joins hermaphroditic duct close to seminal vesicle. Eggs numerous, operculate; developed miracidia with single eye-spot. Vitellarium 2 separated elongate-oval clusters of large follicles, lateral at level

of testis. Excretory pore terminal; details of excretory vesicle not observed (masked by uterus).

Diagnosis

The present material exhibits some of the diagnostic characteristics of, and appears most close morphologically to, *S. obesum*, i.e. bell-shaped concavity at posterior extremity of body, bluntly rounded posterior extremity, long forebody, large pharynx and strongly-developed muscular walls of the genital atrium (Looss, 1902; Fares and Maillard, 1974; Blasco-Costa *et al.* 2009*e*). However, *S. obesum* is characterized by a much larger elongate cylindrical body (BW/BL = 25–34 vs 43–59% (mean 30 vs 48%)), distinctly longer pre-pharynx (185–304 vs 0–76; PL/PHL = 0.8–2.0 vs 0–0.6) and a somewhat smaller sucker width ratio (mean 1 : 0.99 vs 1 : 1.17) due to the ventral sucker being wider than long in *S. brayi* n. sp. Although the measurements of the testis and eggs exhibit overlapping ranges, the upper limits in *S. brayi* are higher for the former (mean 149 × 123 vs 125 × 107 in *S. obesum*) and lower for the latter (mean 50 × 27 vs 53 × 29, see Table 2). Furthermore, the genital atrium appears more prominent in *S. obesum*, whereas it is less muscular in *S. brayi* (mean 82 × 106 vs 67 × 75) and a vesicular *pars prostatica* is apparently absent in the latter species. Finally, both

Table 2. Comparative morphometric data for *Saccocoelium obesum* and *S. brayi* n. sp.

(Standard deviations given for egg-size only.)

Species	<i>S. obesum</i>		<i>S. brayi</i> n. sp.	
	'Large morph 1' of Blasco-Costa <i>et al.</i> (2009 <i>b</i>)			
Morphometric features*	Range	Mean	Range	Mean
Body length	1384–1574	1477	810–997	879
Maximum body width	397–483	440	387–488	428
Oral sucker length	124–144	134	101–142	118
Oral sucker width	137–187	151	99–157	130
Pre-pharynx length	185–304	241	0–76	56
Pharynx length	142–177	160	132–187	150
Pharynx width	104–134	115	85–159	123
Oesophagus length	185–316	269	89–271	167
Ventral sucker length	132–154	140	99–132	116
Ventral sucker width	127–177	149	116–170	147
Hermaphroditic sac length	245–350	295	187–299	241
Hermaphroditic sac width	149–185	167	144–177	164
Genital atrium length	66–101	82	40–83	67
Genital atrium width	96–114	106	58–96	75
Internal seminal vesicle length	127–187	155	99–158	125
Internal seminal vesicle width	53–119	86	63–94	80
External seminal vesicle length	62–109	86	46–114	65
External seminal vesicle width	56–73	67	43–53	50
Testis length	86–154	125	114–210	149
Testis width	80–127	107	94–172	123
Ovary length	78–114	101	53–99	76
Ovary width	71–94	80	81–109	93
Vitelline masses length	88–125	102	71–123	97
Vitelline masses width	46–100	72	54–68	59
Egg length	50–58	53 ± 2	43–55	50 ± 3
Egg width	26–31	29 ± 1	26–29	27 ± 1
Forebody length	511–756	607	314–374	341
Post-uterine field length	127–202	153	76–157	117
Post-caecal field length	349–448	386	230–326	267
Post-testicular field length	147–319	264	73–225	161
Sucker length ratio	1 : 0.99–1.08	1 : 1.04	1 : 0.87–1.09	1 : 0.99
Sucker width ratio	1 : 0.88–1.05	1 : 0.99	1 : 1.00–1.28	1 : 1.17
BW/BL (%)	25–34	30	43–59	48
FO/BL (%)	37–49	41	35–41	38
HSL/VSL (%)	159–261	212	189–227	202
TEND/BL (%)	10–23	18	9–23	18
CEND/BL (%)	24–29	26	26–33	30

* *Abbreviations*: BW/BL (%), maximum body width as a percentage of body length; FO/BL (%), length of the forebody as a percentage of body length; HSL/VSL (%), hermaphroditic sac length as a percentage of ventral sucker length; TEND/BL (%), post-testicular field length as a percentage of body length; CEND/BL (%), post-caecal field length as a percentage of body length.

multivariate approaches (PCA and LDA, see below) clearly indicate a separate allocation of the specimens of *S. obesum* ('large morph 1' from the western Mediterranean), which we consider as *S. obesum* (*sensu stricto*), and specimens of *S. brayi*. The above comparisons coupled with the consistent multivariate morphometric differentiation and observed genetic divergence (see below) support the distinct species status of *S. brayi* n. sp.

Four morphotypes of *S. tensus* ex *Liza* spp.

In addition to the material from *L. aurata* and *L. ramado* identified and described as *S. tensus*

in Blasco-Costa *et al.* (2009*e*), which we label as 'morphotype 1' here, we found 3 isolates in *L. ramado* (labelled as 'morphotypes 2–4') that closely resembled *S. tensus* (Fig. 2A–D). It is worth noting that 'morphotype 1' was collected in fish from the Ebro Delta, 'morphotype 2' was the only species of *Saccocoelium* found in the brackish water lagoon near Santa Pola, whereas 'morphotypes 3 and 4' were collected in hosts fished off Santa Pola.

The specimens tentatively identified as *S. tensus* generally showed a morphological homogeneity, but the 3 morphotypes ex *L. ramado* exhibited gradually increasing upper ranges for most morphometric




Fig. 2. *Saccocoelium tensum* ex *Liza aurata* and *L. ramado*, ventral views with uterus in outline. (A) 'Morphotype 1' ex *L. aurata*. (B) 'Morphotype 2' ex *L. ramado*. (C) 'Morphotype 3' ex *L. ramado*. (D) 'Morphotype 4' ex *L. ramado*. Scale bars = 200 μm .

features (Table 3), the larger 2 forms being distinctly larger and with measurements of the ventral sucker and the length of genital atrium, testis, vitelline and postcaecal fields varying outside the ranges for 'morphotypes 1 and 2'. Specimens of all 3 morphotypes were distinctly more elongate than those of 'morphotype 1', as shown by the lower upper limits (30–39 vs 56%) and means (30–33 vs 42%) for the ratio BW/BL (see also Fig. 2A–D). These differences were reflected in the separation of the specimens in the multivariate analyses (see below). Although the morphometric differentiation in the 2-dimensional plane of the PCA was not as clear for the 'morphotypes 1 and 2', we distinguished the latter prior to sequencing in the following characters: (i) more elongate, narrower body; (ii) long forebody; and (iii) hermaphroditic sac shorter in relation to the size of the ventral sucker, which also appears large in relation to the body (Fig. 2A–B). As shown in Table 3, the specimens of 'morphotype 2' also exhibit larger means for the size of the pharynx, suckers, genital atrium and eggs, and the length of the pre-pharynx, oesophagus and internal seminal vesicle, but possess smaller gonads. The specimens of this morphotype were all gravid adults bearing 30–89 eggs (30, 36, 44, 48, 76, 68 and 89, respectively), and there was no

correlation between the number of eggs and body size ($P=0.119$).

Two Saccocoelium morphotypes ex *M. cephalus*

We did not expect to find yet another form parasitizing *M. cephalus* in the Ebro Delta, an extensively sampled locality in which 2 new species, *S. cephalis* and *S. currani*, were already described (Blasco-Costa *et al.* 2009e; unfortunately, our attempts to obtain sequences for the latter failed). However, we distinguished morphologically a third form in this host, *Saccocoelium* sp., which differed from *S. cephalis* (actually described from the voucher specimens of the present molecular study; see Blasco-Costa *et al.* 2009e) in its distinctly larger and more elongate body (BW/BL = 22–26% vs 26–42%; mean 24 vs 36%) with maximum width at the level of the ventral sucker (vs at the junction of the first and second body thirds), larger suckers, testes, ovary and vitelline masses, and somewhat smaller eggs (Fig. 3A–B; Table 4). The specimens of *Saccocoelium* sp. also possessed a longer oesophagus, more anteriorly terminating caeca (at mid-body vs mid-hindbody) and a more anterior location of the most posterior uterine loops (UEND/BL = 22–42 vs 8–16%). The size differences could not

Table 3. Comparative morphometric data for the four morphotypes of *Saccocoelium tensum* ex *Liza aurata* and *L. ramado*

(Abbreviations as in Table 2.)

Morphotype	1		2		3	4
	<i>L. aurata</i>		<i>L. ramado</i>		<i>L. ramado</i>	<i>L. ramado</i>
Host						
Morphometric features	Range	Mean	Range	Mean	Range	n=1
Body length	853–1133	1011	890–1271	1101	1240–1510	1865
Maximum body width	321–548	428	278–434	356	521–584	560
Oral sucker length	105–139	125	110–148	137	101–152	154
Oral sucker width	114–149	134	139–162	149	139–167	181
Pre-pharynx length	0–63	16	5–45	32	33–99	68
Pharynx length	109–152	121	110–137	128	130–154	152
Pharynx width	94–142	107	110–132	120	137–170	171
Oesophagus length	149–268	218	192–308	235	220–240	309
Ventral sucker length	94–137	117	114–157	131	137–147	177
Ventral sucker width	106–149	135	140–185	156	139–177	180
Hermaphroditic sac length	180–314	230	131–238	211	235–349	316
Hermaphroditic sac width	111–258	149	104–168	143	197–253	233
Genital atrium length	38–51	42	42–76	59	61–114	76
Genital atrium width	51–61	53	50–76	62	43–83	73
Internal seminal vesicle length	61–202	94	101–126	116	134–177	149
Internal seminal vesicle width	35–111	57	45–76	60	86–114	99
External seminal vesicle length	44–147	87	48–87	64	111–170	147
External seminal vesicle width	43–81	62	40–70	54	73–114	94
Testis length	83–167	116	58–134	101	170–235	266
Testis width	67–172	91	57–91	73	106–197	316
Ovary length	66–116	87	46–106	73	81–137	—
Ovary width	51–130	83	53–87	66	116	—
Vitelline masses length	74–131	103	74–151	104	137–159	163
Vitelline masses width	35–68	53	35–67	48	71–73	76
Egg length	37–49	44	45–51	48	43–47 (45)	—
Egg width	21–27	24	24–28	26	25–27 (26)	—
Forebody length	256–390	325	325–456	393	353–458	584
Post-uterine field length	25–250	120	92–227	138	86–129	175
Post-caecal field length	195–407	310	240–378	320	503–663	863
Post-testicular field length	134–380	264	211–336	284	245–539	385
Sucker length ratio	1:0.80–1.10	1:0.94	1:0.85–1.06	1:0.96	1:0.97–1.32	1:1.15
Sucker width ratio	1:0.82–1.13	1:1.01	1:1.03–1.14	1:1.09	1:0.83–1.06	1:0.99
BW/BL (%)	37–56	42	29–38	33	37–39	30
FO/BL (%)	26–36	32	34–40	36	27–30	31
HSL/VSL (%)	158–255	196	105–184	161	201–237	179
TEND/BL (%)	16–36	26	23–31	26	17–36	21
CEND/BL (%)	23–39	31	24–33	29	47	46

be attributed to growth, since some of the specimens of the larger form, *Saccocoelium* sp., were neogravid. However, lack of sufficient numbers of fully-gravid worms prevented confident identification of the species.

Morphometric variation of *Saccocoelium* spp.

The considerable morphological variability encountered in the present collection required a thorough examination to determine the species status of the morphotypes. We, therefore, applied the approach adopted in previous studies on the Haploporinae (see Blasco-Costa *et al.* 2009c, e). First, PCA was applied to assess and visualize the overlap of the morphological characteristics of the

Saccocoelium spp. The first two principal components of the PCA run on the correlation matrix between 13 metrical variables of the 8 morphotypes explained 71.6% of the variation in the data-set comprising 51 specimens. The size of the suckers and the length of the forebody had the highest coefficients on the first component, which explained 56% of the total variance (eigenvalue 7.8), whereas testis size and the length of post-uterine field had important contributions to the second principal component, which explained a further 15.6% of the variance (eigenvalue 2.2). A plot of the specimens in the first plane of the PCA (Fig. 4A) shows, along both the first and second axis, 2 well-separated groups that correspond to the 2 morphotypes from *M. cephalus*. Specimens of *S. obesum* (*sensu stricto*) and *S. brayi* n. sp. appeared




Fig. 3. *Saccocoelium* morphotypes ex *Mugil cephalus*. (A) *S. cephalis*, ventral view of a paratype with uterus in outline. (B) *Saccocoelium* sp., ventral view with uterus in outline. Scale bars = 200 μ m.

close, but separated along the first axis. On the other hand, the morphotypes of *S. tensum* did not show a clear clustering pattern. The specimens of *S. brayi* n. sp. appeared closer and overlapped morphotypes of *S. tensum* in this analysis based on morphometric data only. However, the new species can be readily distinguished from *S. tensum* using only the structure of the posterior extremity of the body.

Secondly, the backward stepwise LDA procedure run on 13 metrical variables separated the 7 morphotypes with 92% accuracy (Fig. 4B) (Wilk's Lambda = 0.00121; approximate $F_{(24, 144)} = 35.2$, $P < 0.0001$). The first canonical function clearly discriminates the two morphotypes ex *M. cephalus* and *S. obesum* (*s. s.*) from the remaining specimens, whereas the second canonical function contributes

to the discrimination between the two morphotypes from *M. cephalus*, between *S. obesum* (*s. s.*) and *S. brayi* n. sp. and between the former and the morphotypes of *S. tensum*. The specimens of *S. obesum* (*s. s.*), 'morphotype 1' of *S. tensum* and the two morphotypes from *M. cephalus* which formed distinct clusters in Fig. 4B were all correctly assigned to their *a priori* groups whereas 2 specimens of *S. brayi* n. sp. were misclassified as 'morphotype 1' of *S. tensum* (accuracy 71%) and 1 specimen of 'morphotype 2' and 'morphotype 3' of *S. tensum* each were misclassified as *S. brayi* (accuracy of 85 and 67%, respectively); these uncertainties are indicated by the higher dispersion and overlap among the samples of *S. tensum* and *S. brayi*. The discriminatory power of the model was associated with

Table 4. Comparative morphometric data for *Saccocoelium cephalis* and *Saccocoelium* sp. ex *Mugil cephalus* (Abbreviations as in Table 2.)

Species	<i>S. cephalis</i>		<i>Saccocoelium</i> sp.	
	Range	Mean	Range	Mean
Morphometric features				
Body length	496–664	583	875–1088	985
Maximum body width	173–230	207	198–270	236
Oral sucker length	64–83	76	90–96	95
Oral sucker width	78–104	88	104–126	113
Pre-pharynx length	0–14	4	0–22	10
Pharynx length	46–58	53	51–58	53
Pharynx width	42–51	47	46–62	55
Oesophagus length	77–192	143	216–288	255
Ventral sucker length	53–66	57	64–75	72
Ventral sucker width	60–69	65	64–80	74
Hermaphroditic sac length	142–166	154	155–200	175
Hermaphroditic sac width	94–104	100	106–147	118
Genital atrium length	32–48	40	37–40	39
Genital atrium width	42–56	49	38–58	47
Internal seminal vesicle length	70–110	94	88–133	107
Internal seminal vesicle width	32–54	43	64–80	73
External seminal vesicle length	48–99	69	54–107	71
External seminal vesicle width	32–37	33	46–85	60
Testis length	77–136	107	218–355	284
Testis width	61–99	80	128–189	156
Ovary length	51–96	61	70–114	92
Ovary width	38–61	52	72–120	93
Vitelline masses length	58–93	76	70–142	116
Vitelline masses width	26–56	40	42–55	47
Egg length	42–43	42	39	—
Egg width	22–23	22	20	—
Forebody length	171–200	187	298–336	317
Post-uterine field length	48–91	64	235–368	313
Post-caecal field length	99–272	207	378–548	475
Post-testicular field length	66–154	124	163–474	311
Sucker length ratio	1:0.64–0.82	1:0.71	1:0.71–0.78	1:0.76
Sucker width ratio	1:0.63–0.84	1:0.74	1:0.60–0.77	1:0.66
BW/BL (%)	26–42	36	22–26	24
FO/BL (%)	28–37	32	31–36	33
HSL/VSL (%)	228–313	273	239–267	248
TEND/BL (%)	13–27	21	19–52	32
CEND/BL (%)	20–44	33	47–56	51

only 4 variables. The length of the testis and the length of the muscular genital atrium exhibited strong correlation with the first axis, and the length of the forebody and the width of the ventral sucker were associated with the discrimination along the second axis of the 2-dimensional plane.

Molecular analysis

Sequences of both 28S and ITS2 rRNA gene regions were obtained from a total of 21 isolates of the 8 morphotypes of *Saccocoelium* spp. Sequences for the 2 rDNA regions from multiple isolates of *S. tensum*, *S. obesum* and *Saccocoelium* sp. (Table 1) showed no variation within the sequences for either region. The 28S and ITS2 sequences for all morphotypes examined were aligned together with the sequences for the two outgroup taxa. The alignment of the 28S rDNA incorporated a total of 1189

included characters (bp and gaps) and the alignment of the ITS2 represented a total of 451 characters. Comparative sequence analysis revealed 4 unique genotypes for each of the two rDNA regions examined: (i) *S. obesum* (*s. s.*); (ii) *S. brayi* n. sp.; (iii) *S. tensum* (all 4 morphotypes); and (iv) *S. cephalis/Saccocoelium* sp.

Divergence in the 28S sequence between species of *Saccocoelium* ranged from 0.8 to 4.3% (9–51 nucleotide sites) and those in the ITS2 sequence ranged from 1.6 to 7.2% (7–33 nt) (Table 5). The smallest divergences were observed between the two morphotypes of *S. obesum* (*sensu lato*) [*i.e.* *S. obesum* (*s. s.*) and *S. brayi* n. sp.], both showing the highest percentage of sequence difference to the other two *Saccocoelium* genotypes. However, the two morphotypes from *M. cephalus*, which appeared morphologically distinct (*S. cephalis* and *Saccocoelium* sp., see above), shared the same genotype for both regions




Fig. 4. Plots of the 52 specimens of *Saccocoelium* spp. (A) In the first plane of the PCA. (B) Against the first and second canonical discriminant functions (LDA). Key to morphotypes of *S. tensum*: 'Morphotype 1', open circles; 'Morphotype 2', light grey circles; 'Morphotype 3', dark grey circles.

Table 5. Pairwise nucleotide sequence comparisons between taxa, calculated as percentage of nucleotide differences (gaps treated as missing data) for the aligned ITS2 (above the diagonal; $N=451$ bps) and 28S rDNA (below the diagonal; $N=1189$ bps) sequences

Taxon	<i>S. cephalii</i>	<i>S. tensum</i>	<i>S. obesum</i>	<i>S. brayi</i>	<i>H. benedeni</i>	<i>F. gibsoni</i>
<i>Saccocoelium cephalii</i>	—	3.1	7.2	7.2	6.3	16.8
<i>Saccocoelium tensum</i>	2.5	—	7.2	6.3	6.5	16.0
<i>Saccocoelium obesum</i>	4.1	3.4	—	1.6	8.8	17.5
<i>Saccocoelium brayi</i> n. sp.	4.3	3.7	0.8	—	9.0	17.4
<i>Haploporus benedeni</i>	7.7	7.1	7.7	8.0	—	15.4
<i>Forticulcita gibsoni</i>	9.4	9.1	9.1	9.6	9.7	—




Fig. 5. Tree topology derived from the combined 28S and ITS2 rRNA gene sequences using Bayesian analysis with posterior probability values above and MP bootstrap values below the branches. *Abbreviations*: i followed by number, sequence isolate number; LA, *Liza aurata*; LR, *Liza ramado*.

analysed and the 4 morphotypes of *S. tensum* (1 from *L. aurata* and 3 from *L. ramado*) were genetically identical for the 2 rDNA regions. The Mantel test, aimed to assess the congruence between the genetic (ITS2 region) and morphometric differentiation between the 7 morphotypes, resulted in a high *P*-value ($P=0.32$), thus confirming the lack of correlation between the genetic and morphometric distance matrices in the present material.

Figure 5 presents the phylogram generated with BI analyses of the combined dataset (28S and ITS2). MP of the combined dataset produced a single most-parsimonious tree (length 336, consistency index 0.896) with strong nodal support for all clades (not shown). The tree topologies, obtained in independent analyses of the 2 gene regions, shared the same branching patterns and showed similar levels of support. Within the ingroup, 2 strongly supported clades were recognized, one formed by morphotypes of *S. tensum* from *L. aurata* and *L. ramado* and

morphotypes from *M. cephalus*, and the other comprising *S. obesum* (*s. s.*) and *S. brayi* n. sp., the latter being subtended by a longer branch.

DISCUSSION

Our study is the first parallel molecular and morphological attempt focused on characterization of a group of congeneric species within the Haploporidae, a poorly known digenean family characterized by a long history of inadequate descriptions, poor specific diagnoses, scattered records and extensive synonymy (Overstreet and Curran, 2005). The family has also been found to be highly labile in its placement in the most comprehensive molecular phylogenetic analysis of the Digenea performed to date (Olson *et al.* 2003), and this reflects the difficulty in characterizing the group at effectively all taxonomic levels. Combining sequence data analysis with a detailed morphological and

multivariate morphometric study of the specimens has allowed the demonstration of cryptic diversity among the closely related species of *Saccocoelium*. One important result is that molecular data corroborate our decisions based on morphology (Blasco-Costa *et al.* 2009*e*) with respect to the distinct status of 3 species, *S. obesum* (*s. s.*), *S. tensum* and *S. cephalis*, thus rejecting the hypothesis of a single species infecting sympatric mullets in the Mediterranean (e.g. Dawes, 1947; Mikailov, 1958; Fischthal and Kunz, 1963; Ferreti and Paggi, 1965; Moravec and Libosvářský, 1975). Furthermore, the analysis of both gene regions supported the recognition of another cryptic species, *S. brayi* n. sp. Species recognition based on sequence divergence in the ITS2 in cases when low interspecific variation is detected between congeners is difficult; one is doomed to either fail to recognize multiple cryptic species or provide 'species' destined for synonymy (see Nolan and Cribb, 2005 for a review). The amount of morphological and genetic variation between *S. obesum* (*s. s.*) and *S. brayi* n. sp. was lower than that observed between *S. tensum* and *S. cephalis*, and this would suggest their recent separation. In addition, the morphological differences between the morphotypes of *S. obesum* (*s. l.*) were confirmed by the multivariate morphometric analyses. These, coupled with the observed sequence divergence in the 28S and ITS2 regions, the former slightly above the minima (e.g. 0.2–0.4% in the Cryptogonimidae, see Miller and Cribb, 2007*a, b*) and the latter higher than or closely approaching the lowest levels reported between congeneric taxa in other marine digenean systems (e.g. 0.5% in the Didymozoidae, see Anderson and Barker, 1998; 0.3% in the Sanguinicolidae, see Nolan and Cribb, 2006; 0.4–1.4% in the Cryptogonimidae, see Miller and Cribb, 2007*a, b*), support our decision to recognize the distinct species status of *S. brayi* n. sp. Further sequencing, especially of material from the Black Sea, will be useful to test our hypothesis for higher diversity in the *S. obesum* (*s. l.*) complex (Blasco-Costa *et al.* 2009*e*) and to reveal the divergence rates within this morphologically and genetically distinct lineage of *Saccocoelium*.

The lack of genetic differentiation among the 4 morphotypes of *S. tensum* is in accord with our initial hypothesis of a single morphologically variable species based on the long list of synonyms and intraspecific variation detected in descriptions from various mullet hosts (Blasco-Costa *et al.* 2009*e*), and the lack of qualitative differentiating features and the gradually overlapping ranges for the metrical data in the present material. However, the finding that *Saccocoelium* sp. isolates had identical sequences to *S. cephalis*, especially of the more variable ITS2 region, was unexpected in view of the apparent boundaries indicated by comparative morphology and multivariate statistical analysis; this is reflected in the larger number of isolates sequenced. One

explanation for the observed lack of genetic differentiation in the ITS2 region within the *S. tensum* – *S. cephalis* clade may be a slower rate of evolution than that found in the *S. obesum* (*s. s.*) + *S. brayi* n. sp. lineage, the latter exhibiting a considerably longer branch length. If this is the case, it is possible that the ITS2 region may not provide enough resolution to guarantee distinguishing between all combinations of closely related haploporid species (see also Nolan and Cribb, 2005). Nevertheless, ensuing from the fact that to date just a single convincing example for identical ITS2 sequences in genuinely different species exists within the Digenea (Blair *et al.* 1997; discussed in Nolan and Cribb, 2005), we adopted a conservative approach, considering the distinct species status only for the *Saccocoelium* isolates that are supported by both morphological and molecular evidence. However, we provide sufficient data for their morphological differentiation and have deposited voucher material of the morphotypes studied here, in order to enable their recognition should future studies on different loci (e.g. ITS1 or mitochondrial genes) offer evidence validating their species distinction.

Divergence of the two clades of *Saccocoelium* appears to be associated with a change in the intermediate host group rather than with diversification related to the definitive hosts. Life-cycle data available for *S. tensum* and *S. obesum* (if extended to the other members of their respective clades) tend to support this suggestion. Cercariae of *S. tensum* develop in the hydrobiid snails *Hydrobia acuta* and *H. ventrosa*, whereas those of *S. obesum* develop in the rissoid snails *Rissoa* spp. (Fares and Maillard, 1974; Deblock, 1980; the former authors also reported numerous unsuccessful attempts to infect experimentally *Hydrobia* spp. with *S. obesum*). The free-living cercariae of both species encyst in the open by attaching to the substrate and are thus available for passive ingestion by the definitive pump-filtering, detritivorous (Cardona *et al.* 2001) mullet hosts in which the adult stages develop. Host-parasite data, although recently updated, are still wanting due to the large body of non-documented records (i.e. with no supportive evidence for the species identification). Nevertheless, documented records for both *S. obesum* and *S. tensum* include 5 of the 6 widespread sympatric Mediterranean mullet species (i.e. *Chelon labrosus*, *Mugil cephalus*, *Liza aurata*, *L. ramado* and *L. saliens*) as definitive hosts (see Blasco-Costa *et al.* 2009*e* for details). This fact and the current lineage division suggest that parasite specificity at the level of the definitive host may not serve as an important force for speciation in the mullet-haploporid system studied. This situation is similar to that observed in a Mediterranean sparid-digenean system comprising 3 host species which occur in sympatry (Jousson *et al.* 2000). A wide open encounter filter (*sensu* Combes, 1995; Combes and

Théron, 2000, Poulin, 2006) in our system resulting from the combination of the distinctly 'passive' transmission of the infective metacercariae which are ingested non-selectively and the habitat/trophic overlap between the mullet host species would tend to reduce the possibility of co-existence of genetically isolated mullet-specific lineages of *Saccocoelium*.

However, the higher than previously thought species diversity, confirmed by the present molecular data at the limited geographical scale of the study (c. 500 km, with parasite species co-occurring in the same locality) and the coexistence of divergent parasite lineages in the same host species indicate the action of factors linked to the haploporid life-cycle which modify the effect of the enhanced encounter with the definitive hosts. Two non-exclusive hypotheses can be suggested for the observed patterns of species and genetic diversity in our system. The first, and an obvious prediction, is that the existence of more species of *Saccocoelium* reflects adaptation to different sympatric snail hosts. The relationship with the snail intermediate hosts can be viewed as the most important component of the compatibility filter (*sensu* Combes, 1995, Poulin, 2006), since digeneans exhibit the highest specificity to their molluscan hosts (Pearson, 1972; Adamson and Caira, 1994), and this supports our suggestion. Estimation of both hydrobiid and rissoid species diversity in the Mediterranean is difficult due to problems in species delimitation (Anistratenko and Stadnichenko, 1994; Wilke and Davis, 2000 and references therein). However, a particularly rich set of sibling species was established in European rissoaceans (Russo and Patti, 2005 and references therein) and members of *Hydrobia* were shown to represent morphostatic species radiations (i.e. '... considerable, rapid speciation[s] with low anatomical diversification'; see Davis, 1994; Wilke and Davis, 2000). It is therefore possible that speciation within *Saccocoelium* is associated with cryptic diversification of the first intermediate rissoid and hydrobiid hosts. Sequencing of both snail hosts and larval stages in the western Mediterranean lagoons would provide a test of this hypothesis.

Our second hypothesis is that the higher parasite diversity is a result of local adaptation governed by larval dispersal of the snail intermediate hosts. Larval spatial distributions and dispersal ability have been linked to genetic differentiation among free-living marine organisms (e.g. Tatarenkov and Johannesson, 1998; Boisselier-Dubayle and Gofas, 1999; Riginos and Victor, 2001). Of particular relevance to the system under study is the fact that *H. ventrosa* (first intermediate host of *S. tensus* and 2 other haploporid species) is a species with a direct development (i.e. crawl-away juveniles emerge after metamorphosis from egg masses deposited on the substrate), and this results in high population level differentiation and low gene flow between

populations (Foltz, 2003; Wilke and Davis, 2000). The poor dispersal and the heterogeneity of the habitats (see Bartoli and Gibson, 2007 for comment on lagoonal types in the western Mediterranean) may therefore provide a setting for the development of differential susceptibility in the populations of this host towards infections with *Saccocoelium* spp. and haploporids in general. Thus, the possibility of speciation *via* local adaptation, resulting from the spatial structure of the first intermediate host populations, might be high in haploporids.

Summarizing the results of our study, we conclude that distinct species status is only valid for the *Saccocoelium* isolates that are supported by both morphological and molecular evidence. Our data suggest that delimiting species using solely morphological criteria may be misleading; however, we do not rule out the possibility for even higher species diversity within the studied digenean group. By describing sequence and morphological divergence across the lowest taxonomic levels, we provide a test case that demonstrates which genetic and morphological markers can be used for diagnostic analysis in the Haploporidae.

We are grateful to David Gibson, Natural History Museum, London, for advice in the course of the study and to Tomas Scholz, Institute of Parasitology, Biology Centre of the Academy of Sciences of the Czech Republic, for comments on the manuscript. This research was supported by INTAS (J.A.B., A.K., grant 03-51-5998); Ministry of Science and Education of Spain (J.A.B., grant CGL2008-02701); FP6-MTKD (J.A.R., A.K., grant CT-2004-003175); and Institute of Parasitology (A.K., grants Z60220518 and LC522). I.B.-C. benefits from a doctoral fellowship from the Ministry of Science and Education of Spain (AP-2004-2999) and a visiting studentship to the Natural History Museum in London.

REFERENCES

- Adamson, M. L. and Caira, J. N.** (1994). Evolutionary factors influencing the nature of parasite specificity. *Parasitology* **109**, S85–S95.
- Anderson, G. R. and Barker, S. C.** (1993). Species differentiation in the Didymozoidae (Digenea): Restriction fragment length differences in internal transcribed spacer and 5.8S ribosomal DNA. *International Journal for Parasitology* **23**, 133–136.
- Anderson, G. R. and Barker, S. C.** (1998). Inference of phylogeny and taxonomy within the Didymozoidae (Digenea) from the second internal transcribed spacer (ITS2) of ribosomal DNA. *Systematic Parasitology* **41**, 87–94.
- Anistratenko, V. V. and Stadnichenko, A. P.** (1994). *Fauna Ukrainyi*, Tom 29. Mollyuski, Vyipusk 1, Kniga 2: Littorinoobraznyie, Rissoiobraznyie. Naukova Dumka, Kiev.
- Bartoli, P. and Gibson, D. I.** (2007). Synopsis of the life cycles of Digenea (Platyhelminthes) from lagoons of the northern coast of the western Mediterranean. *Journal of Natural History* **41**, 1553–1570.

- Bickford, D., Lohman, D. J., Sohdi, N. S., Ng, P. K. L., Meier, R., Winker, K., Ingram, K. K. and Das, I.** (2007). Cryptic species as a window on diversity and conservation. *Trends in Ecology and Evolution* **22**, 148–155.
- Blair, D., Agatsuma, T., Watanobe, T., Okamoto, M. and Ito, A.** (1997). Geographical genetic structure within the human lung fluke, *Paragonimus westermani*, detected from DNA sequences. *Parasitology* **115**, 411–417.
- Blasco-Costa, I., Balbuena, J. A., Kostadinova, A. and Olson, P. D.** (2009a). Interrelationships of the Haploporinae (Digenea: Haploporidae): A molecular test of the taxonomic framework based on morphology. *Parasitology International* **58**, 265–269.
- Blasco-Costa, I., Gibson, D. I., Balbuena, J. A., Raga, J. A. and Kostadinova, A.** (2009b). A revision of the Haploporinae Nicoll, 1914 (Digenea: Haploporidae) from mullets (Mugilidae): *Haploporus* Looss, 1902 and *Lecithobotrys* Looss, 1902. *Systematic Parasitology* **73**, 107–133.
- Blasco-Costa, I., Montero, F. E., Balbuena, J. A., Raga, J. A. and Kostadinova, A.** (2009c). A revision of the Haploporinae Nicoll, 1914 (Digenea: Haploporidae) from mullets (Mugilidae): *Dicrogaster* Looss, 1902 and *Forticulcita* Overstreet, 1982. *Systematic Parasitology* **72**, 187–206.
- Blasco-Costa, I., Montero, F. E., Gibson, D. I., Balbuena, J. A. and Kostadinova, A.** (2009d). A revision of the Haploporinae Nicoll, 1914 (Digenea: Haploporidae) from mullets (Mugilidae): Two new haploporine genera and a key to the genera of the subfamily. *Systematic Parasitology* **72**, 207–215.
- Blasco-Costa, I., Montero, F. E., Gibson, D. I., Balbuena, J. A., Raga, J. A. and Kostadinova, A.** (2009e). A revision of the Haploporinae Nicoll, 1914 (Digenea: Haploporidae) from mullets (Mugilidae): *Saccocoelium* Looss, 1902. *Systematic Parasitology* **72**, 159–186.
- Boisselier-Dubayle, M. C. and Gofas, S.** (1999). Genetic relationships between marine and marginal-marine populations of *Cerithium* species from the Mediterranean Sea. *Marine Biology* **135**, 671–682.
- Cardona, L., Royo, P. and Torras, X.** (2001). Effects of leaping grey mullet *Liza saliens* (Osteichthyes, Mugilidae) in the macrophyte beds of oligohaline Mediterranean coastal lagoons. *Hydrobiologia* **462**, 233–240.
- Chambers, C. B. and Cribb, T. H.** (2006). Phylogeny, evolution and biogeography of the Quadrifoliovariinae Yamaguti, 1965 (Digenea: Lecithasteridae). *Systematic Parasitology* **63**, 61–82.
- Combes, C.** (1995). *Interactions Durables. Écologie et Evolution du Parasitisme*. Masson, Paris, France.
- Combes, C. and Théron, A.** (2000). Metazoan parasites and resource heterogeneity: constraints and benefits. *International Journal for Parasitology* **30**, 299–304.
- Davis G. M.** (1994). Molecular genetics and taxonomic discrimination. *The Nautilus Supplement* **2**, 3–23.
- Dawes, B.** (1947). *The Trematoda of British Fishes*. Ray Society, London, UK.
- Deblock, S.** (1980). Inventaire des Trématodes larvaires parasites des Mollusques Hydrobia (Prosobranches) des côtes de France. *Parassitologia* **22**, 1–105.
- Donald, K. M., Kennedy, M., Poulin, R. and Spencer H. G.** (2004). Host specificity and molecular phylogeny of larval Digenea isolated from New Zealand and Australian topshells (Gastropoda: Trochidae). *International Journal for Parasitology* **34**, 557–568.
- Edgar, R. C.** (2004). MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **5**, 113.
- Fares, A. and Maillard, C.** (1974). Recherches sur quelques Haploporidae (Trematoda) parasites des Muges de Méditerranée Occidentale: systématique et cycles évolutifs. *Zeitschrift für Parasitenkunde* **45**, 11–43.
- Ferretti, G. and Paggi, L.** (1965). Ridescrizione di *Saccocoelium obesum* Looss, 1902 (sin. *Saccocoelium tensum* Looss, 1902) trematoda parassita di *Mugil cephalus*. *Rivista Parassitologia* **26**, 229–239.
- Fischthal, J. H. and Kuntz, R. E.** (1963). Trematode parasites of fishes from Egypt. Part V. Annotated record of some previously described forms. *Journal of Parasitology* **49**, 91–98.
- Foltz, D. W.** (2003). Invertebrate species with nonpelagic larvae have elevated levels of nonsynonymous substitutions and reduced nucleotide diversities. *Journal of Molecular Evolution* **57**, 607–612.
- Hall, T. A.** (1999). Bioedit: a user-friendly biological sequence editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium*, Ser. 41, 95–98.
- Holzer, A. S., Sommerville, C. and Wootten, R.** (2004). Molecular relationships and phylogeny in a community of myxosporeans and actinosporeans based on their 18S rDNA sequences. *International Journal for Parasitology* **34**, 1099–1111.
- Huelsenbeck, J. P., Ronquist, F., Nielsen, R. and Bollback, J. P.** (2001). Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* **294**, 2310–2314.
- Jousson, O. and Bartoli, P.** (2001). Molecules, morphology and morphometrics of *Cainocreadium labracis* and *Cainocreadium dentecis* n. sp. (Digenea: Opecoelidae) parasitic in marine fishes. *International Journal for Parasitology* **31**, 706–714.
- Jousson, O., Bartoli, P. and Pawlowski, J.** (2000). Cryptic speciation among intestinal parasites (Trematoda: Digenea) infecting sympatric host fishes (Sparidae). *Journal of Evolutionary Biology* **13**, 778–785.
- Leung, T. L. F., Keeney, D. B. and Poulin, R.** (2009). Cryptic species complexes in manipulative echinostomatid trematodes: when two become six. *Parasitology* **136**, 241–252.
- Littlewood, D. T. J., Curini-Galletti, M., Herniou, E. A.** (2000). The interrelationships of Proseriata (Platyhelminthes: Seriata) tested with molecules and morphology. *Molecular Phylogenetics and Evolution* **16**, 449–466.
- Looss, D. A.** (1902). Die Distomen-Unterfamilie der Haploporinae. *Archives de Parasitologie* **6**, 129–143.
- Lockyer, A. E., Olson, P. D. and Littlewood, D. T. J.** (2003). Utility of complete large and small subunit rRNA genes in resolving the phylogeny of the Neodermata (Platyhelminthes): implications and a review of the cercomer theory. *Biological Journal of the Linnean Society (London)* **78**, 155–171.
- Maddison, D. R. and Maddison, W. P.** (2005). *MacClade 4: Analysis of Phylogeny and Character*

- Evolution. Version 4.08*. Sinauer Associates, Sunderland, MA, USA.
- Manly, B. F. J.** (1997). *Randomisation, Bootstrap and Monte Carlo Methods in Biology*, 2nd Edn. Chapman and Hall, London, UK.
- Mikhailov, T. K.** (1958). Parasite fauna of *Mugil saliens* Risso of the Caspian Sea. *Zoologicheskyy Zhurnal* **37**, 373–378 (In Russian).
- Miller, T. L. and Cribb, T. H.** (2007a). Two new cryptogonimid genera (Digenea, Cryptogonimidae) from *Lutjanus bohar* (Perciformes, Lutjanidae): analyses of ribosomal DNA reveals wide geographic distribution and presence of cryptic species. *Acta Parasitologica* **52**, 104–113.
- Miller, T. L. and Cribb, T. H.** (2007b). Coevolution of *Retrovarium* n. gen. (Digenea: Cryptogonimidae) in Lutjanidae and Haemulidae (Perciformes) in the Indo-West Pacific. *International Journal for Parasitology* **37**, 1023–1045.
- Miura, O., Kuris, A. M., Torchin, M. E., Hechinger, R. F., Dunham, E. J. and Chiba, S.** (2005). Molecular-genetic analyses reveal cryptic species of trematodes in the intertidal gastropod, *Batillaria cumingi* (Crosse). *International Journal for Parasitology* **35**, 793–801.
- Moravec, F. and Libosvářský, J.** (1975). Intestinal helminths from grey mullets, *Mugil capito* Cuvier and *M. cephalus* L., of Lake Borullus, A.R.E. *Folia Parasitologica* **22**, 279–281.
- Nolan, M. J. and Cribb, T. H.** (2004). Two new blood flukes (Digenea: Sanguinicolidae) from Epinephelinae (Perciformes: Serranidae) of the Pacific Ocean. *Parasitology International* **53**, 327–335.
- Nolan, M. J. and Cribb, T. H.** (2005). The use and implications of ribosomal DNA sequencing for the discrimination of digenean species. *Advances in Parasitology* **60**, 101–163.
- Nolan, M. J. and Cribb, T. H.** (2006). *Cardicola* Short, 1953 and *Braya* n. gen. (Digenea: Sanguinicolidae) from five families of tropical Indo-Pacific fishes. *Zootaxa* **1265**, 1–80.
- Olson P. D. and Tkach V. V.** (2005). Advances and trends in the molecular systematics of the parasitic platyhelminthes. *Advances in Parasitology* **60**, 165–243.
- Olson, P. D., Cribb, T. H., Tkach, V. V., Bray, R. A. and Littlewood, D. T. J.** (2003). Phylogeny and classification of the Digenea (Platyhelminthes: Trematoda). *International Journal for Parasitology* **33**, 733–755.
- Overstreet, R. M. and Curran, S. S.** (2005). Family Haploporidae Nicoll, 1914. In *Keys to the Trematoda*. Vol. 2 (ed. Jones, A., Bray, R. A. and Gibson, D. I.), pp. 129–165. CAB International, Wallingford, UK.
- Pearson, J. C.** (1972). A phylogeny of life-cycle patterns of the Digenea. *Advances in Parasitology* **58**, 760–788.
- Posada, D. and Crandall, K. A.** (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**, 817–818.
- Poulin, R.** (2006). *Evolutionary Ecology of Parasites*, 2nd Edn. Princeton University Press, Princeton, NJ, USA.
- Riginos, C. and Victor, B. C.** (2001). Larval spatial distributions and other early life-history characteristics predict genetic differentiation in eastern Pacific blennioid fishes. *Proceedings of the Royal Society London, B* **268**, 1931–1936.
- Russo, G. F. and Patti, F. P.** (2005). Early life history of two closely related gastropods, *Rissoa auriscalpium* and *Rissoa italiensis* (Caenogastropoda: Rissoidae). *Marine Biology* **147**, 429–437.
- Ronquist F. and Huelsenbeck, J. P.** (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- Swofford, D. L.** (2002). *PAUP – Phylogenetic Analysis Using Parsimony. Ver. 4*. [Computer software and manual]. Sinauer Associates, Sunderland, MA, USA.
- Tatarenkov, A. and Johannesson, K.** (1998). Evidence of a reproductive barrier between two forms of the marine periwinkle *Littorina fabalis* (Gastropoda). *Biological Journal of the Linnean Society (London)* **63**, 349–365.
- Tkach, V. V., Grabda-Kazubska, B., Pawlowski, J. and Swiderski, Z.** (1999). Molecular and morphological evidences for close phylogenetic affinities of the genera *Macrodera*, *Leptophallus*, *Metaleptophallus* and *Paralepoderma* (Digenea, Plagiorchioidea). *Acta Parasitologica* **44**, 170–179.
- Wilke, T. and Davis, G. M.** (2000). Intraspecific mitochondrial sequence diversity in *Hydrobia ulvae* and *Hydrobia ventrosa* (Hydrobiidae: Rissoidae: Gastropoda): Do their different life histories affect biogeographic patterns and gene flow? *Biological Journal of the Linnean Society (London)* **70**, 89–105.